

PTE
12/25/72

Terraclor® SOIL FUNGICIDE 40% DUST

ACTIVE INGREDIENT:

Pentachloronitrobenzene

40%

INERT INGREDIENTS:

Total

60%
100%**CAUTION
KEEP OUT OF REACH OF CHILDREN**

If swallowed: Do not induce vomiting. Do not get product in eyes. May cause skin irritation. Wash thoroughly with soap and water after use of eye contact lenses or if irritation develops. Wash thoroughly after use of gloves. Wear gloves when handling the powder or dust. Avoid breathing dust or fume stuffs.

USE CAUTIONS

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not plant any root crop not registered for PCNB in rotation on PCNB treated soil.

Dispose of empty bags by burning (KEEP OUT OF SMOKE)

NOTICE TO BUYER

The manufacturer or distributor of this product makes no claim as to its effectiveness for any purpose other than that set forth in the directions for use. The manufacturer or distributor shall not be liable for any damages resulting from the use of this product in accordance with other methods or procedures than those recommended. It is the responsibility of the user to determine the suitability of this product for his particular use.

DIRECTIONS FOR USE**PEANUTS (Southern Stem and Root Rot — Sclerotium rolfsii):**

Surface Band Treatment: Apply 25 lbs. per acre (per 12,500 linear feet of row) in a 12 inch band centered on the row, at pegging time. Where cultural practices permit, this may be mixed and applied simultaneous with land-plaster.

Pre-plant soil mix: Apply 25 lbs. per 12,500 linear feet of row in an 8 to 12 inch band centered on the row and mix to a depth of 2 to 4 inches. Application may be made simultaneous with planting, using suitable application equipment.

Soil Mix During Cultivation: Apply in split applications during three cultivations at the rate of 8 lbs. per 12,500 linear feet of row per application. Attach two delivery tubes per row ahead of inside cultivator sweeps (one on each side) and adjust so as to treat a band of 12 inches wide centered on the row. Do not pile untreated soil on crown of plant.

NOTE: Do not feed treated vines to livestock. Do not allow hogging down of treated peanuts.

TOMATOES AND PEPPERS (Southern Blight — Sclerotium rolfsii):

Trench Dust: Dust 18 pounds per 7,300 linear feet for flat tomatoes, or 18 per 10,900 linear feet for staked tomatoes, and for 14,500 linear feet of row for peppers, over open "V" trench just before setting transplants. Place in bottom of trench then press walls of trench against stems of young transplants. Note: When cultivating hot peppers, do not let plants get too tall as they may fall over and damage the stems of plants.

EPA Reg. No. 1200-27

EF-10-76

Terr'allor

SOIL FUNGICIDE

ACTIVE INGREDIENT:

Pentachloronitrobenzene

40%

INERT INGREDIENTS:

Total

60%

100%

PEANUTS (Southern Stem and Root Rot - *Sclerotium rolfsii*)

Surface Band Treatment: Apply 25 lbs. per acre (per 12,500 linear feet of row) in a 12 inch band centered on the row at pegging time. Where cultural practices permit, the band may be band applied simultaneously with lime-plaster.

Pre-plant soil mix: Apply 25 lbs. per 12,500 linear feet of row in a 12 inch band centered on the row at the time of tillage or 2 weeks after. Application may be made at any time during plowing, harrowing, or other soil preparation.

Soil Mix During Cultivation: Apply 25 lbs. per 12,500 linear feet of row in a 12 inch band centered on the row at the time of tillage or 2 weeks after. After cultivation, the band may be band applied simultaneously with lime-plaster.

NOTE:

The product is a water-soluble powder. It is recommended that it be applied dry. If it becomes wet, it should be dried before application.

For PCNB in

soil treatment, add 1 lb. of PCNB to 100 lbs. of Terr'allor.

Trend. Dose.

See label for details.