

# OLCO

## 46% EMULSIFIABLE CHLORDANE

Contains 4# Technical Chlordane per gallon

For control of Chiggers, Ticks, Chinch Bugs, Cut Worms, Fire Ants, Pharaoh Ants, Grasshoppers, Wire Worms, White Grubs, Japanese Beetle larvae, Asiatic Beetle Larvae.

ACTIVE INGREDIENTS ..... 90%

Technical Chlordane\*..... 46%

Petroleum Oil ..... 44%

INERT INGREDIENTS ..... 10%

\*Equivalent to 27.6% Octachloro-4, 7 methanotetrahydroindane and 18.4% related compounds.

EMULSIFIES READILY WITH WATER

### CAUTION: KEEP OUT OF REACH OF CHILDREN

CAUTION: Chlordane is toxic and can be absorbed through the skin. Harmful if swallowed. Avoid inhaling spray mists. Avoid contamination of feed and food stuffs.

ANTIDOTES: External - In case of spillage on skin, wash with soap and water. Internal - Drink emetic of mustard and call a physician.


.....Gallons

Compounded by

# OLIVER CHEMICAL CO.

CINCINNATI, OHIO

USDA Registration No. 1275-13



### DIRECTIONS

Lawn Applications. The following insects are controlled by the recommended dosages as given in terms of volume of 46% Emulsifiable Chlordane to be added to 5 gallons of water and sprayed or sprinkled on 1,000 square feet of lawn area. Be careful to not contaminate oriental fish ponds. Birds feeding in treated areas may be harmed.

For control of CHIGGERS & TICKS, mix 3 tablespoons in 5 gallons of water - After spray is applied do not sprinkle lawn for 48 hours.

For control of CHINCH BUGS & CUT WORMS, mix 6 tablespoons in 5 gallons of water. After spray is applied sprinkle lawn thoroughly.

For control of FIRE ANTS, PHAROAH ANTS and other species commonly found in lawns, mix 2 1/4 tablespoons in 5 gallons of water. After spray is applied do not sprinkle lawn for 48 hours. Apply spray thoroughly in and around ant hills.

For control of GRASSHOPPERS, mix 2 1/4 tablespoons in 5 gallons of water. After spray is applied do not sprinkle lawn for 48 hours.

For control of WHITE GRUBS, JAPANESE BEETLE LARVE, ASIATIC LARVAE, 1/2 to 1 measuring cup (4 to 8 fl. ozs.) in 5 gallons of water. For solid type soils, as peat and mucks, use 1 measuring cup of 46% Emulsifiable Chlordane, for clay and loams use 3/4 cup, and for sandy soil use 1/2 cup. After applying spray, it is important to soak lawn thoroughly with additional water.

For control of WIREWORMS, mix 3 to 12 tablespoons in 5 gallons of water - the amount of 46% Chlordane to use depends on the type of soil. Sandy soils require least, clays and loams require 4 to 6 tablespoons, peat and mucks require 9 to 12 tablespoons. After spray is applied, it is important to thoroughly soak the soil with additional water.

For control of WIREWORMS in TRANSPLANT WATER (feeder barrel). To protect tobacco seedlings against wire worms add 1/2 pint of 46% Emulsifiable Chlordane to 50 gallon barrel of transplant water. Apply at the rate of 6 to 8 barrels per acre.