

DIRECTIONS FOR USE

COMPLETE COVERAGE SPRAY

Ground equipment recommendations call for 30-50 gallons of water per acre. Airplane recommendations call for five gallons of water per acre.

FRUITS

APPLES — Use 2 to 3 quarts Staples Parathion per acre in 30-50 gallons of water for the control of aphid, red-banded leaf roller, codling moth, scale insects and leaf miner starting at pre-pink stage and after petal fall. If infestation is not controlled, continue at 10-14 day intervals as needed. Do not apply to McIntosh type apples. The allowable dosage that may be applied is 3 gallons per acre (6.0 lbs. parathion). Do not apply within 14 days of harvest.

EBERRIES — for the control of maggots and thrips, use at the rate of 2 pints per acre in 30-50 gallons water. Allowable dosage is 2½ pints per acre (0.6 lbs. parathion). Do not apply within 14 days of harvest.

PEACHES — Use 2 quarts per acre in 30-50 gallons of water for the control of plum curculio, oriental fruit moth, scale insects, cat-facing insects, green peach aphid and fruit tree leaf roller.

FOR AREAS OTHER THAN CALIFORNIA: Full coverage spray control of scale insects — Do not apply within 14 days of harvest. No not apply more than 5 lbs. actual parathion per acre (2½ gallons Staples Parathion per year). The allowable dosage is 2 gallons per acre (4.0 lbs. parathion).

FOR CALIFORNIA: Do not apply within 21 days of harvest. Do not apply more than once after bloom. Do not apply more than 2½ lbs. actual parathion (5 quarts of this brand of parathion) per acre between January 1 and Harvest. The allowable dosage is 5 pints per acre (2.5 lbs. parathion).

PUMPS AND PRUNES — Use 2 quarts per acre in 30-50 gallons of water for the control of plum curculio, spider mite aphid, leaf roller, leafhopper and bud moth. Do not apply within 14 days of harvest. The allowable dosage is 2 gallons per acre (4 lbs. parathion).

STRAWBERRIES — Use 1 quart per acre in 30-50 gallons of water for the control of leaf roller and red spider mite, making applications at 7-10 day intervals as required. Do not apply within 14 days of harvest. The allowable dosage is 1½ quarts (0.8 lbs. parathion) per acre.

VEGETABLES

BEANS — Use 1 quart per acre in 30-50 gallons of water for control of Mexican bean beetle, leaf-hopper, aphid, red spider, armyworm, leaf roller and leaf miner. Apply at 7 to 10 day intervals as required. Do not apply within 15 days of harvest. The allowable dosage is 1½ quarts per acre (0.8 lbs. parathion).

BEETS — Use 1 quart per acre in 30-50 gallons of water for the control of webworm and aphid. Do not apply within 15 days of harvest (21 days if tops are to be used for food or feed). The allowable dosage of this brand of parathion is 3 pints per acre (0.8 lbs. parathion).

BEST DOCUMENT AVAILABLE

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

25% PARATHION

CONTAINS TWO POUNDS PARATHION PER GALLON

25% PARATHION forms an emulsion when diluted with water to any desired strength. It will not clog fine orifices and is suitable for use in all power operated sprayers and aircraft sprayers.

ACTIVE INGREDIENTS

| | |
|---|-------|
| Parathion (O, O-diethyl, O-p-nitrophenyl thiophosphate) | 25.0% |
| Aromatic Petroleum Hydrocarbon | 67.0% |
| INERT INGREDIENTS | 8.0% |

TOTAL.....100.0%

EPA Registration No. 1345-8

EPA EST. NO. 01345 ME 01

DANGER

- Keep Out of Reach of Children

☠ POISON ☠

See Antidote Statement and Other Required Warning

Statements on the Back Panel

Control No.

NET CONTENTS GALLONS

Manufactured By

C. W. Staples, Inc.

PRESQUE ISLE, MAINE

BEST DOCUMENT AVAILABLE

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

25% PARATHION

CONTAINS TWO POUNDS PARATHION PER GALLON

25% PARATHION forms an emulsion when diluted with water to any desired strength. It will not clog fine orifices and is suitable for use in all power operated sprayers and aircraft sprayers.

| | |
|---|---------------|
| ACTIVE INGREDIENTS | |
| Parathion (O, O-diethyl, O-p-nitrophenyl thiophosphate) | 25.0% |
| Aromatic Petroleum Hydrocarbon | 67.0% |
| INERT INGREDIENTS | |
| | 8.0% |
| TOTAL | 100.0% |

EPA Registration No. 1345-8
EPA EST. NO. 01345 ME 01

DANGER

- Keep Out of Reach of Children

POISON

See Antidote Statement and Other Required Warning

Statements on the Back Panel

Control No.

NET CONTENTS GALLONS

Manufactured By

C. W. Staples, Inc.

PRESQUE ISLE, MAINE

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE Con't.

SUGAR BEETS — Use this brand of parathion at the rate of 1 quart per acre in 30-50 gallons of water for sugar beet webworm. Apply when insects first appear and repeat at 7 day intervals as necessary. Do not apply within 15 days of harvest (21 days if the tops are used for food or feed). The allowable dosage is 3 pints (0.8 lbs. parathion) per acre.

CARROTS — Use ¾ pints per acre in 30-50 gallons of water for control of aphid. Do not use within 15 days of harvest. The allowable dosage is 2 quarts (1 lb. parathion) per acre.

SQUASH AND MELONS — For the control of cucumber beetle, melon worm, serpentine leaf miner, aphid, and pickle-worm, use 1 quart per acre in 30-50 gallons of water. Do not apply to squash within 15 days of harvest. Do not apply to melons within 7 days of harvest. The allowable dosage is 1 quart (0.5 lbs. parathion) per acre.

PEAS — For the control of aphid, use 1 quart per acre in 30-50 gallons of water. Do not apply within 10 days of harvest. Where pea vines are used for forage, do not apply within 15 days of harvest. The allowable dosage is 1 quart per acre (0.5 lbs. parathion).

POTATOES — Use 1 quart of this brand of parathion per acre in 30-50 gallons of water for control of flea beetles, leaf hopper and aphid. Do not apply within 5 days of harvest. The allowable dosage is 2 quarts per acre (1.0 lb. of parathion).

TOMATOES — For tomatoes raised as field crops, use 1 quart per acre in 30-50 gallons of water for control of flea beetle, leafhopper and aphid. Do not apply within 10 days of harvest. Allowable dosage is 2 quarts (1.0 lb. parathion) per acre.

CORN — For the control of European corn borer use 2 quarts per acre in 30-50 gallons of water. Apply first spray when 50% of eggs are in the blackhead stage. Make three applications 5 to 7 days apart. Do not apply within 12 days of harvest or cutting for forage. The allowable dosage is 2 quarts (1.0 lb. parathion) per acre.

CABBAGE — Use 1 pint per acre in 30-50 gallons of water for the control of diamond-back moth, imported cabbage worm, army worm, aphid and thrip. Do not apply within 7 days of harvest in which case only the allowable dosage is 1 pint per acre (0.25 lbs. parathion). Do not apply within 10 days of harvest in which case the allowable dosage is 1 quart per acre (0.5 lb. parathion).

CUCUMBERS — For the control of cucumber beetle, serpentine leaf miner, aphid and pickle-worm, use 1 quart per acre in 30-50 gallons of water. Do not apply before plants begin to vine. Do not apply within 15 days of harvest. The allowable dosage is 1 quart (0.5 lbs. parathion) per acre.

FORAGE CROPS — For the control of aphid, grasshoppers, army worm found on wheat, oats, barley, clover, grass, hay and alfalfa use 1 quart per acre in 30-50 gallons of water. Do not apply within 15 days of cutting. The allowable dosage is 1½ quarts per acre (0.8 lbs. parathion) for alfalfa, grass and hay. For wheat, oats, barley and clover, the allowable dosage is 3 quarts (1.5 lbs. parathion) per acre. **NOTE ON SPRAYING PASTURES WITH PARATHION.** Do not apply within 15 days of pasturing.

BEST DOCUMENT AVAILABLE

STAF ES 25% PARATHION

W A R N I N G

Rapidly absorbed through the skin.

Poisonous if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin or on clothing. Keep away from heat or open flame. Wear a mask or respirator of a type approved by the U. S. Department of Agriculture for parathion protection.

o - - - - o - - - - o - - - - o

"This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label." "Do not reuse empty containers. Destroy containers when empty."

o - - - - o - - - - o - - - - o

Wash hands and face thoroughly with soap and water before eating or smoking. Wear protective clothing, rubber gloves and goggles. Keep all unprotected persons out of treated area or vicinity where there may be danger of drift. Treated areas should not be reentered until volatile residues have dissipated.

ANTIDOTE STATEMENT

SYMPTOMS OF POISONING — A sense of "tightness" in chest; stomach pains; vomiting and diarrhea.

ANTIDOTE — Internal: Give a teaspoonfull of salt in a glass of warm water and repeat until vomit fluid is clear. Have patient lie down and keep quiet. Call a physician at once.

ANTIDOTE — External: If the skin is contaminated, wash immediately with soap and warm water.

TO PHYSICIAN — Compound exerts muscarine-like effects, that is, stimulation of the parasympathetic nervous system. A parasympathetic depressant, such as atropine, is indicated to relieve symptoms. The administration of oxygen is helpful.

NOTICE

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

"DO NOT USE, POUR, OR SPILL OR STORE NEAR HEAT OR OPEN FLAME."

BEST DOCUMENT AVAILABLE

0 3 1 3 0

0 3 1 3 0