

NATIONAL

APOLLO 445

ACCEPTED

MAY 24 1978

U.S. Dept. of Federal Insecticide,
Fungicide and Rodenticide Act,
as amended for the pesticide
registration system
EPA Reg. No. 1769-143

National Chemsearch APOLLO 445 is specifically formulated for controlling broadleaf weeds in ornamental turf grasses as well as Wild Onion and Wild Garlic. National Chemsearch APOLLO 445 controls surface creeping weeds as well as broadleaf weeds. May be used on Merion, Bermuda, Kentucky Blue Grass, Fescues and ryegrasses, on lawns, rights-of-way, golf course fairways, and other non-crop commercial turf areas.

Weeds Controlled:

- | | |
|---------------------|--------------|
| Buckhorn | Pigweed |
| Clover | Plantain |
| Common Chickweed | Purslane |
| Curled Dock | Ragweed |
| Dandelion | Red Clover |
| English Daisy | Red Sorrel |
| Ground Ivy | Stitchwort |
| Knotweed | Speedwell |
| Lambsquarters | White Clover |
| Mallow | Wild Garlic |
| Mouse-ear Chickweed | Wild Onion |
| Morning Glory | Yarrow |

DIRECTIONS FOR USE

Mix 6 ounces with 2 to 4 gallons of water and apply uniformly to 1000 square feet. Application can be made by using a sprayer or watering can. Repeat application after 3 to 4 weeks if necessary.

Either before or after application, it is wise to properly fertilize the lawn so as to encourage the lawn to heal as rapidly as possible, filling in the bare areas where the weeds have died away. Once the weeds have been removed, the bare areas should be reseeded. Bare areas may be reseeded 3 to 4 weeks after application.

ACTIVE INGREDIENTS:

- * Diethanolamine salt of 2 (2 Methyl 4-chlorophenoxy) propionic acid 8.0%
- ** Diethanolamine salt of 2, 4 Dichlorophenoxyacetic acid 8.0%

INERT INGREDIENTS: 84.0%

- * Equivalent to 5.7% 2 (2 Methyl 4 chlorophenoxy) propionic acid
- ** Equivalent to 5.7% 2, 4 Dichlorophenoxyacetic acid.

CAUTION:

KEEP OUT OF REACH OF CHILDREN
(See right panel for additional cautions)

EPA Reg. No. 1769-143-AA

NET CONTENT

NATIONAL CHEMSEARCH

2727 Chemsearch Blvd. Irving, Texas 75062

NOT INTENDED FOR HOUSEHOLD
USE OR RESALE

WHEN TO APPLY: Apply in spring or fall when weeds are actively growing. Knotweed is controlled more easily in spring or early summer when weeds are young.

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes or clothing. Avoid inhaling spray mist. Avoid contamination of water intended for irrigation or domestic purposes. Do not reuse shipping containers; destroy by crushing and burying where possible. Avoid spray drift to susceptible plants as this product may injure grapes, tomatoes, beans, peas, other vegetables, legumes, ornamentals, etc. (Coarse sprays are less likely to drift.) Because of danger of contamination, do not use same spray equipment for applying insecticides, etc. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not transport with or store near seeds, fertilizers, insecticides or fungicides. Do not use in or around greenhouses. Store at temperatures above 32°F(0°C.) If allowed to freeze, thaw and re-mix before using. **KEEP OUT OF REACH OF CHILDREN.**

CAUTION: Do not apply when turf is suffering from drought conditions or during very hot weather or damage to grass may result. Do not mow for several days before or after application. Withhold sprinkling for 24 hours after application. Do not apply during periods of heavy rainfall. Do not use on newly seeded areas. Do not use if you want clover in your turf. In some areas bent grasses, carpet, buffalo and St. Augustine grasses may be injured; therefore do not use on turf where these grasses predominate.

Local conditions may affect the use of this chemical. Consult Agricultural Experiment Station or Extension Service weed specialists for specific recommendations as to dosage and time of application and for control of other weeds.

DO NOT STORE WITH FOODSTUFFS

EPA Est. No. 4140

78

CA 1 IN 1 NJ 1 PR 1 TX 1

2 78