

ACCEPTED

9-29-70

1812-91

Commercial



10% SEVIN

with sulphur



ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-Methylcarbamate)	-----	10%
Sulphur	-----	60%
INERT INGREDIENTS	-----	30%
Total	-----	100%

Sulphur Fineness 93 - passing No. 325 USS Sieve

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Do not contaminate feed stuffs, feed, feeding troughs or receptacles.

DIRECTIONS: Use sufficient amount of dust to obtain full coverage. Apply when insects or their signs appear. Repeat at 5 to 7 day intervals or as necessary for good control. Use in accordance with label directions. Sevin may be applied up to time of harvest.

COTTON: To control Boll Weevil, Boll Worms, Thrips, Flea Beetles, Fall Army Worms, Stink Bugs, dust at rate of 15-20 lbs. per acre according to infestation.

BEANS: Mexican Bean Beetle. Dust at rate of 5 lbs. per acre according to infestation.

PEANUTS: Velvet Bean Caterpillar, Peanut Worms, the Fall Army Worm and Thrips. Apply from 7-10 lbs. per acre depending upon size of plant and infestation. Repeat as often as needed.

For protection of Honey Bees, do not use if bees are visiting plants when honey bees are visiting the crop. When necessary to use during the season, beekeepers will advance to locate hives at a safe distance until one week after application.

U.S. PAT. NO. 2,812,001

Parramore & Griffin

Valdosta, Ga.