



2% PHOSDRIN DUST

ACTIVE INGREDIENTS:

Alpha isomer of 2-Carbomethoxy-1-methylvinyl Dimethyl Phosphate 1.2%
 Related compounds3%

INERT INGREDIENTS:

..... 98.0%
 Total 100.0%



POISON

*Equivalent to 2.0%w Phosdrin Insecticide

DANGER: KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Do not swallow or get in eyes, on skin, or clothing. Wear clean natural rubber gloves, protective clothing and goggles. Wear mask or respirator of a type passed by the U. S. Bureau of Mines for Phosdrin Insecticide protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into eyes, immediately flush with water for at least ten minutes and get medical attention. In case of spillage on person or clothing, immediately remove all contaminated clothing and wash skin with soap and water. Wash clothing before reuse.

Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas must not be entered until drifting insecticide and volatile residues have been dissipated.

Do not contaminate food or feed products.

Do not reuse container. Bags should be burned or buried. Stay well away from smoke.

This product is toxic to fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label.

NOTE TO PHYSICIAN: ATROPINE IS ANTIDOTAL.

DIRECTIONS: Make first application when insects appear. Repeat applications as often as necessary to maintain control.


Cabbage, Mustard, Turnips, Summer Squash. Against Aphids, Cabbage Loopers, leafhoppers, mites, climbing cutworms and grasshoppers. Use 25 lbs. per acre Do not apply within 1 day of harvest on Squash and Cabbage. Do not apply within 3 days of harvest on Mustard and Turnips.

Beans. Against Aphids, grasshoppers, leafhoppers, mites, mexican bean beetles. Use 25 lbs. per acre. Do not apply within 1 day of harvest.

Peas. Against Aphids, grasshoppers, leafhoppers, mites, climbing cutworms. Use 25 lbs. per acre. Do not apply within 1 day of harvest.

Collards. Against Aphids, cabbage loopers, grasshoppers, mites, climbing cutworms. Use 25 lbs. per acre. Do not apply within 3 days of harvest.

Cucumbers. Against Aphids, grasshoppers, leafhoppers, mites. Use 25 lbs. per acre. Do not apply within 1 day of harvest.



PARRAMORE & GRIFFIN

E.P.A. No. 1812-104
 E.P.A. No. 1812-104

Valdosta, Georgia