




# 5% SEVIN - COPPER - SULPHUR

## ● ● PEANUT DUST ● ●



### ACTIVE INGREDIENTS

Carbaryl (1-naphthyl N-methylcarbamate) .....	5.0%
Copper, as metallic (in Copper Oxide) .....	4.0%
Sulphur .....	71.4%
INERT INGREDIENTS: .....	19.6%
Total .....	100.0%

Carbaryl is the registered trademark of ICI-Naphthyl N-Methylcarbamate Sulphur fineness: 100 mesh, 100% U.S. Sieve

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**

### CAUTION

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Do not contaminate foodstuffs, feed, feeding troughs or receptacles.

**DIRECTIONS:** Use sufficient amount of dust to obtain full coverage. Apply when insects or their signs appear. Repeat at 5 to 7 day intervals or as necessary for good control. Use in accordance with label directions. Sevin may be applied up to time of harvest. General Suggestion: Apply 30 lbs. per acre, repeated as often as needed. At least every 14 days.

Sulphur may burn foliage when temperatures are excessively high.

### PEANUTS

5% Sevin-Copper-Sulphur is for use on peanuts for the control of Leaf Spot, Fall Army Worms & Thrips. Consult your County Agent or Experiment Station for complete details on the peanut dusting program best suited to your local conditions.

General Suggestion: Apply 30-35 lbs. per acre, repeated as often as needed. At least every 14 days.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

PARRAMORE & GRIFFIN

1970, Rev. 11-1810-197

Valdosta, Georgia