

DIRECTIONS

LICE ON BEEF CATTLE, HOGS, SHEEP AND GOATS: Mix 1 gallon **MALATHION 57%** Concentrate in 100 gallons water. Spray completely animals, pens and litter. Repeat only as needed. Do not treat Dairy Goats.

TICKS ON BEEF CATTLE, SHEEP AND GOATS: Mix 1 gallon **MALATHION 57%** Concentrate in 100 gallons water. Spray animals completely. Repeat in 2 weeks if needed. Do not treat Dairy Goats.

HORN FLIES ON BEEF CATTLE: Mix 1 to 1½ gallons **MALATHION 57%** Concentrate in 100 gallons water. Spray completely. Repeat at two week intervals if needed.

BACK RUBBERS: For reduction of Lice on Beef Cattle: Treat back rubbers with a mixture of 1 pint of **MALATHION 57%** Concentrate in 4 gallons of diesel fuel oil. Do not use crankcase or lubrication oil. Repeat every 2-3 weeks. May also be a reduction in horn flies. Do not use on dairy cattle.

POULTRY (Chickens, Ducks, Geese, Turkeys): Direct application for Northern Fowl Mite, Poultry Lice and as a supplement to Premise Treatment for Chicken Red Mites: Mix 1 ounce (2 tablespoonfuls) of **MALATHION 57%** Concentrate per gallon of water and spray at the rate of 1 gallon per 100-150 birds. Repeat in 4 to 6 weeks or when necessary.

POULTRY HOUSES: Northern Fowl Mite, Chicken Red Mite, poultry Lice and Flies: Mix 2 ounces (4 tablespoonfuls) of **MALATHION 57%** in 1 gallon of water. Spray liberally walls, ceilings, roosts and nests, forcing spray into all cracks and crevices.

POULTRY TICKS: Mix 6-7 ounces of **MALATHION 57%** Concentrate per gallon of water. Spray liberally on walls, ceilings, and adjacent areas, forcing spray into all cracks and crevices. Repeat as needed.

DOGS AND CATS: Fleas: Spray with a mixture of 1 ounce (2 tablespoonfuls) of **MALATHION 57%** Concentrate per gallon of water. Wet animals thoroughly. Repeat as needed, but not more than once a week. Do not treat kittens and puppies under four weeks of age.

ANIMAL QUARTERS: Fleas: Remove manure or debris. Spray kennels, pens, yards, lawns and under houses with a mixture of 5 ounces of **MALATHION 57%** per gallon of water at the rate of 1 gallon per 1,000 square feet. Use a garden sprayer.

HOUSEHOLD INSECTS: Roaches, Ants, Spiders, Silverfish, Crickets, Clover Mites, Earwigs in and around Homes, Dairies and Food Processing Plants. Dilute 1 ounce (2 tablespoonfuls) of **MALATHION 57%** in one quart of water (4 ounces per gallon.) Spray or paint around window frames, baseboards, drainboards, under sinks, stoves, cracks and crevices, and other areas frequented by insects. Do not contaminate food or utensils. Repeat application as necessary. **CAUTION:** This product may cause staining.

EPA REG. NO. 1926-73

EPA EST NO.

ACCEPTED

SEP 18 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 1926-73

With Corn



NAVY TOX PREMIUM GRADE

Emulsifiable Concentrate to be used as a grain protectant and for the control of Lice, Ticks, Hornflies on Beef Cattle, Lice on Hogs, Lice, Mites, Ticks on Poultry. For use as a fly bait. Flies and Fleas on Dogs and Cats. Certain Household insects. Can be used in Back Rubbers and Stock Oilers.

ACTIVE INGREDIENTS		90.09%
Malathion*	57.00%	
Aromatic Petroleum Derivative Solvent	33.09%	
INERT INGREDIENTS		9.91%
*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.		

**CAUTION: KEEP OUT OF REACH OF CHILDREN
DO NOT USE, POUR, SPILL, OR STORE NEAR
HEAT OR OPEN FLAME**

See side panel for further precautions.

SOLD BY

navy brand.

MALATHION 57% CONCENTRATE

FLY CONTROL: Remove lactating dairy animals from building before treating. As a bait spray use 2 quarts **MALATHION 57%** Concentrate with 1 quart corn syrup or molasses or 5 pounds of sugar in 25 gallons of water. Apply around foundations and entrances of dwellings; in floor areas, runways, feed rooms in dairy barns and out buildings, also to walls and other fly-frequented areas. Use sprinkling can or sprayer. Also spray manure piles and other debris or fly breeding areas. Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed. **DO NOT APPLY THIS DOSAGE TO ANIMALS.** Remove calves under one month of age before treating. Do not use in milk processing rooms. Maintain sanitary conditions around barns, outbuildings and other areas to prevent fly breeding. Repeat applications as necessary.

**CAUTION:
KEEP OUT OF REACH OF CHILDREN
AND DOMESTIC ANIMALS**

Harmful if swallowed or absorbed through skin. Avoid breathing mist. Avoid contact with skin, eyes, clothing. Wash thoroughly after using.

MALATHION 57% may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid.
NOTE TO PHYSICIAN: Atropine is antidotal.

Do not apply to lactating dairy cattle. Do not apply to dry dairy cattle within 2 weeks of freshening. Do not treat animals under 1 month of age. Do not use on dairy goats. After spraying, swine should be kept out of sun and wind for a few hours.

Avoid contamination of feed, food stuffs, food containers and water troughs.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

Food utensils such as teaspoons or tablespoons should not be used for food purposes after use with pesticides.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective Information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE OF WARRANTY

navy brand makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statements on this label.

Net Contents

11.5 Gal

ACCEPTED WITH COMMENTS