

Controls broadleaved plants, narrow leaved plants, thistle, henbit, wild carrots, common chickweed, mouse-eared chickweed, dandelion, oxalis, spotted spurge, and legumes such as white clover and black medick.

Apply 7 fluid ounces (4 tablespoons) in one gallon of water or more per each 1,000 square feet of lawn surface.

For the control of poison ivy and ground ivy, mix 4 to 8 fluid ounces per 3 gallons of water and spray to wet all foliage thoroughly.

USE DIRECTIONS: Best results are obtained when applied in cooler weather during spring or fall. If used in very hot weather, may cause excessive damage to turf. Do not apply to newly sown turf, nor to stoloniferous grasses such as bent, Bermuda or St. Augustine. Do not use same spray equipment for other purposes, unless cleaned with a suitable chemical cleaner.

CAUTION: Do not take internally. Avoid breathing spray mist. May cause skin irritation. Avoid contact with eyes, skin and clothing. In case of contact flush with plenty of water, wash with soap and water.

Vapors from this product may injure susceptible plants in the immediate vicinity. Do not apply directly to or allow solution or spray drift to contact desirable plants (including but not limited to vegetable, grapes, fruit trees and ornamentals). Coarse sprays are less likely to drift than fine mist sprays.

DISCLAIMER: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herein.

U.S.D.A. Reg. No. 1996-146

Net Contents 8 Ounces

18 3322



SILVEX

LAWN WEED KILLER

ACCEPTED

ACTIVE INGREDIENT

Isooctyl Ester of Silvex (2,2,4,5-Tetrachlorophenoxy propionic acid) 37.4%

INERT INGREDIENTS

Inert 62.6%
Total 100.0%
Equivalent to 26.4% by weight of Silvex, or not less than two pounds of Silvex per gallon.

CAUTION