



NACO
7 1/2% SEVIN* DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)	7.5%
INERT INGREDIENTS	92.5%
TOTAL	100.0%

U. S. Patent Nos. 2,903,478; 3,009,855
*Trademark of Union Carbide Corp. for 1-naphthyl N-methylcarbamate

(18-point) CAUTION!
(12-point) KEEP OUT OF REACH OF CHILDREN
(See side panels for additional cautions and directions)

HARMFUL IF INHALED OR SWALLOWED. DO NOT TAKE INTERNALLY. DO NOT BREATHE DUST.
SKIN CONTACT MAY BE HARMFUL. AVOID CONTACT. Wash thoroughly after using. Take shower or bath after work. Wear regular long sleeved clothing. Change to clean clothing daily.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs and water receptacles.

For protection of honeybees avoid use if possible when honeybees are visiting the crop or neighboring crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance (beyond bee flight range) until one week after application.

DIRECTIONS FOR USE

Read Entire Label. Use in strict accordance with label directions and cautions. Apply when wind is calm at the rates indicated herein. Be sure to cover both sides of the leaves with a light even coat of dust. Repeat application if rains occur within 24 hours after application. Do not graze treated fields or feed trash to dairy or meat animals.

COTTON: To control Boll Weevil, Bollworm, Pink Bollworm, Cotton Aphids, Cotton Leafworm, Lygus Bugs, Cotton Leaf Perforator, Thrips and Cotton Fleahopper apply at the rate of 12 to 25 pounds per acre. Consult your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with this label cautions and directions.

Usually, for early season control, 2 to 3 applications should be made at 7 to 10 day intervals at the rate of 10 to 15 pounds per acre. For mid-season control, apply 15 to 20 pounds per acre. Repeat application at 5 to 7 day intervals until infestation is brought under control. Usually 4 applications will be necessary. Repeat a series of 4 applications at 5 to 7 day intervals if boll weevils re-appear in sufficient numbers to cause damage. For late season control, apply at the rate of 20 to 25 pounds per acre at 5 to 7 day intervals. Anytime bollworms are a threat, when as many as 4 to 5 small worms and eggs are found per 100 terminals examined, apply at least 20 pounds of dust per acre. Direct the dust downward into the tops of the plant where the worms are hatching. Repeat applications at 5 day intervals until infestation is controlled. Where mite infestations are encountered, use a recommended miticide.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

W. R. GRACE & CO.
AGRICULTURAL PRODUCTS DIVISION
BALTIMORE, MARYLAND
NET WEIGHT POUNDS