


NACO
EVALUATED DUST



ACTIVE INGREDIENTS:		
Maneb (Mangencese ethylene bisdithiocarbamate)	4.8	
INERT INGREDIENTS	95.2	
	TOTAL	100.0

(18-point) CAUTION!
 (12-point) KEEP OUT OF REACH OF CHILDREN
 (See side panels for additional cautions)


HARMFUL IF TAKEN INTERNALLY. AVOID INHALATION. AVOID CONTACT WITH SKIN AND EYES.
 In case of skin contact wash with plenty of soap and water. For eyes, flush with water for fifteen minutes and get medical attention.
 In case of accident induce vomiting and call a physician.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Apply this product only as specified on this label. Disposal of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by burying with wastes, or by burning (Keep out of smoke).

GENERAL DIRECTIONS

Apply this product with adequate dusting equipment when the wind is still at the rates specified below. The average amount to use to each application is dependent on the size of the plant, but be sure to cover thoroughly both sides of the leaves with a light even coat of dust.

TOMATOES: To control Early blight, late blight, Septoria leaf spot, Gray leaf spot (Sterphylium) and Anthracnose apply at the rate of 15 - 40 lbs. per acre. Start application when fruit of first cluster is well formed; repeat at 7 to 10 day intervals. (Do not use on tender young seedlings in green houses as injury may occur).



When weather conditions favor disease, more frequent applications may be necessary, starting when plants are set in the field. No time limitations.

POTATOES: To control Early blight and Late blight apply at the rate of 15 - 40 lbs. per acre. Start applications as soon as the plants are 2 to 6 inches high and continue at 5 - 10 day intervals as long as blight threatens. Under severe disease conditions, use the shorter interval or the higher dosages. No time limitations.

CUCUMBERS: To control Downy mildew and Anthracoese apply at the rate of 15 - 40 lbs. per acre. Start treatments as soon as plants begin to run or when disease threatens and continue at 7 to 10 day intervals as long as necessary. Under severe conditions shorten intervals to 3 - 5 days. No time limitations.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

W. R. GRACE & CO.
AGRICULTURAL PRODUCTS DIVISION
BALTIMORE, MARYLAND
NET WEIGHT POUNDS

4-3-68
2124-304