



NACO

PARATHION-SEVIN\* 1-5 DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) .....	5%
Parathion (O,O-diethyl O-p-nitrophenyl thiophosphate) .....	1%
INERT INGREDIENTS .....	94%
TOTAL	<u>100%</u>

U. S. Patent Nos. 2,903,478; 3,009,855

\*Trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate

(18-point)  
(12-point)

WARNING!

KEEP OUT OF REACH OF CHILDREN

(See side panels for additional warnings and directions)

MAY BE FATAL IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN! RAPIDLY ABSORBED THROUGH SKIN! DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING. WEAR NATURAL RUBBER GLOVES, PROTECTIVE CLOTHING AND GOGGLES. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

ANTIDOTES: If swallowed. Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a Physician Immediately!

If on skin. In case of contact remove contaminated clothing and immediately wash skin with soap and water.

This product is toxic to fish and wildlife. Keep out of lakes, streams, and ponds. Apply this product only as specified on this label. Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by burying with wastes.

#### DIRECTIONS FOR USE

READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL. For control of major Tobacco Insects. Hand Dusting Application for Budworms and Tobacco Flea Beetles, when hand dust applications are made on individual plants, apply 6 to 8 pounds per acre. Heavier applications may cause plant injury. Rotary dusters are recommended. Broadcast Application for Hornworms, Budworms, Aphids, Flea Beetles, Suckfly, Grasshoppers and Japanese Beetles. Apply 20 to 40 pounds per acre depending on size of the plants. Use the highest rate only on large plants. Full coverage is essential. Begin treatment when worms are small and repeat as needed. Heavier applications may cause plant injury.

NOTE: If late applications are necessary, treat immediately after priming. Do not apply within 5 days of priming or 15 days of cutting. Avoid getting wet with plant juices when cutting the crop.

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

2 2 07  
1 1 1 44

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

W. R. GRACE & CO.  
AGRICULTURAL PRODUCTS DIVISION  
BALTIMORE, MARYLAND

NET WEIGHT    POUNDS

124-344