


ACCEPTED

NOV 20 1972
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 100-100-100
TO ATTACHED COMMENTS

2124-326



GENERAL DIRECTIONS

Naco Malathion-Methoxychlor-2-2 E.C. may be used in conventional hydraulic sprayers, low-volume ground applicators, and aircraft applicators (use not less than 2 gals. per acre with aircraft applications). Mix this product with water by agitation; agitate the mixture while spraying. If allowed to stand, re-agitate before use.

APPLICATION DIRECTIONS

ALFALFA - To control Alfalfa weevil larvae apply 5 pints (in approximately 25 gallons of water) per acre. Make application in spring when most of the tips are beginning to show feeding. A second application may be necessary. If harvest infestation is severe, treat stubble as soon as crop is removed. To control Clover leaf weevil apply 2 quarts (in approximately 25 gallons of water) per acre. Apply in spring when alfalfa is 2 to 6 inches tall.

To control Alfalfa caterpillar apply up to 2 quarts (in approximately 25 gallons of water) per acre. Apply if there are 10 or more caterpillars per sweep of net, usually when alfalfa is 1/2 grown.

MALATHION-METHOXYCHLOR-2-2 E.C.

ACTIVE INGREDIENTS:

Malathion	22.73%
Methoxychlor	22.73%
Xylene	44.82%

INERT INGREDIENTS

TOTAL	9.72%
	100.00%

0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate
Equivalent to 20.00% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 2.73% other isomers and reaction products.



APPLICATION DIRECTIONS, continued

To control meadow spittlebug apply 2 quarts (in approximately 25 gallons of water) per acre. Apply within a week after eggs hatch.

To control potato leafhopper apply 1 quart (in approximately 25 gallons of water) per acre. Apply when alfalfa is about half grown or earlier if insects become abundant. Do not apply within 7 days of harvest. Do not apply during full bloom.

CLOVER To control clover leaf weevil apply 2 quarts (in approximately 25 gallons of water) per acre. Apply in spring when clover is 2 to 6 inches tall. To control meadow spittlebug apply 2 quarts (in approximately 25 gallons of water) per acre. Apply in spring within a week after eggs begin to hatch. As a fall treatment, apply 3 quarts (in approximately 25 gallons of water) per acre. Make application in early September.

Treat clover in bloom only when bees and other pollinators are not visiting plants. Do not apply closer than 7 days to harvest.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given here-with.

CAUTION! KEEP OUT OF REACH OF CHILDREN

(See side panels for additional cautions)

HARMFUL IF SWALLOWED. AVOID PROLONGED OR REPEATED BREATHING OF SPRAY MIST. AVOID PROLONGED OR REPEATED CONTACT WITH SKIN.

Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. If swallowed, induce vomiting and get medical attention.

This product is toxic to fish. Wildlife in treated areas may be harmed. Keep out of lakes, streams, and ponds. Apply this product only as specified on this label. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

W. R. GRACE & CO.
AGRICULTURAL CHEMICALS
MEMPHIS, TENNESSEE

Amarillo, Tex. • Baltimore, Md. • Cary, N.C. • Charleston, S.C. • Columbus, Ohio • Ft. Pierce, Fla. • Henry, Ill. • Joplin, Mo. • Lansing, Mich. • Nashville, Tenn. • New Albany, Ind. • Perry, Iowa • Portland, Oregon

NET CONTENTS GALLONS

