

NACO

NACO

PARATHION-FERBAM 1-11.4 DUST

PRECAUTIONS

MAY BE FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN! Do not get in eyes, on skin, or on clothing. May cause irritation of eyes, nose, throat and skin. Wear natural rubber gloves, protective clothing and goggles. Workers should use rubber gloves when handling treated plants. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for parathion protection. Keep all unprotected persons out of treated areas of plants, where there may be danger from dust. Do not use in areas where children play. Do not use in areas where people work. Do not use in areas where people live. Do not use in areas where people work. Do not use in areas where people live. Do not use in areas where people work. Do not use in areas where people live.

The product is a restricted use pesticide. It is not to be used by anyone other than a person who has been specifically trained and licensed by the State of California.

NOTICE: Follow directions carefully. Varying method of application, weather & crop conditions, mixtures with other chemicals not specifically recommended & other influencing factors in the use of this product are the condition of the applicator. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given here-with.

ACTIVE INGREDIENTS:

Parathion (O, O-diethyl O-p-nitrophenyl thiophosphate)	1.0%
Ferbam (Ferric dimethyl dithiocarbamate)	11.4%
INERT INGREDIENTS	87.6%

TOTAL 100.0%

DANGER!
KEEP OUT OF REACH OF CHILDREN

SEE HIDE PANEL FOR A COMPLETE LIST OF CROPS



DIRECTIONS FOR USE

1. Apply the required amount of Naco Parathion-Ferbam 1-11.4 Dust uniformly to the foliage with suitable ground dusting equipment.

Tobacco - For use in tobacco plant beds for control of white grub, dust ground thoroughly when plants are dry at the rate of 4 to 6 lbs. per 100 sq. ft. Repeat if necessary.

2. Control of aphid on tobacco plants in early stages of growth at the rate of 2 lbs. per 100 sq. ft. Repeat 4 to 5 days if necessary. Dust at 4 to 5 days after the first application if necessary.

3. Control of beet beet on tobacco plants in early stages of growth at the rate of 2 lbs. per 100 sq. ft. Repeat 4 to 5 days if necessary. Dust at 4 to 5 days after the first application if necessary.

4. Control of cutworm on tobacco plants in early stages of growth at the rate of 2 lbs. per 100 sq. ft. Repeat 4 to 5 days if necessary. Dust at 4 to 5 days after the first application if necessary.

W. R. GRACE & CO.

NET WEIGHT POUNDS

ACCEPTED WITH COMMENTS

ACCEPTED
Feb 7 1970