



NACO

20% TOXAPHENE DUST

ACTIVE INGREDIENT:

Toxaphene (Technical Chlorinated Camphene -Chlorine
Content 67 - 69%)

20%

INERT INGREDIENTS

80%

TOTAL

100%

(18-point)
(12-point)

CAUTION!
KEEP OUT OF REACH OF CHILDREN
(See additional cautions and directions below)

HARMFUL IF SWALLOWED. AVOID CONTACT WITH SKIN, EYES, AND
CLOTHING. AVOID INHALATION OF DUST. Wash thoroughly after use.
Avoid contamination of feed and foodstuffs.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds.
Apply this product only as specified on this label. Dispose of wastes by burying
in non-crop lands away from water supplies. Containers should be disposed by
burying with wastes, or by burning (Keep out of smoke).

DIRECTIONS FOR USE

Apply required amounts of Naco 20% Toxaphene uniformly to the foliage with suitable
ground or airplane equipment when the wind is still. Repeat applications if rain
occurs within 24 hours after applying. Consult your State Agricultural Experiment
Station or your State Agricultural Extension Service for additional information
consistent with this label cautions and directions.

11 - 7 - 72

2124-578

COTTON: To control boll weevil apply 10 - 20 lbs. per acre. Apply on foliage every 3 - 7 days until controlled.

To control boll worm apply 10 - 20 lbs. per acre on foliage every 5 - 7 days until controlled.

To control cotton fleahopper apply 7.5 - 20 lbs. per acre. Apply on foliage as needed.

To control cutworms apply 10 - 20 lbs. per acre. Apply on lower portion of plants and soil surface around plants as needed.

To control fall armyworm apply 10 - 20 lbs. per acre. Apply on foliage as needed.

To control flea beetles apply 10 - 15 lbs. per acre. Apply on foliage as needed.

To control grasshoppers apply 10 - 20 lbs. per acre. Apply on foliage as needed.

Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.

CABBAGE: To control cutworms (soil) apply 10 lbs. per acre broadcast to soil surface as a single preplant treatment or to be applied before the edible parts start to form.

Do not apply within 7 days of harvest if cabbage will be marketed after outer leaves have been removed. Do not apply after heads start to form if cabbage will be marketed with outer leaves intact.

PEANUTS: To control climbing cutworms apply 7.5 - 10 lbs. per acre. Apply as soon as damage is noted.

To control fall armyworm apply 10 lbs. per acre when larvae are young.

ACCEPTED
11-7-72

To control velvetbean caterpillar apply 7.5 lbs. per acre. ²¹²⁴⁻⁵⁷⁸ Apply a few days after eggs hatch or as soon as feeding is noted.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

SOYBEANS: To control bean leaf beetle, armyworm, green cloverworm, and velvetbean caterpillar apply 7.5 lbs. per acre.

Do not apply within 21 days of harvest. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

TOMATOES: To control flea beetles and hornworms apply 15 pounds per acre.

Apply on foliage as needed. Do not apply closer than 3 days of harvest.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

W. R. GRACE & CO.
AGRICULTURAL CHEMICALS
MEMPHIS, TENNESSEE

NET WEIGHT POUNDS