


NACO

NACO



2124-624  
ACCEPTED  
Dec 8 1971

# SEVIN-SULPHUR 10-50 DUST

USDA Reg. No. 2124-624

### DIRECTIONS FOR USE

Apply when the wind is calm at the rate indicated herein. Be sure to cover both sides of the leaves with a light even coat of dust. Repeat application if rains occur within 24 hours after application.

**BEANS**—To control Mexican bean beetles and leafhoppers apply 10 lbs. per acre. To control corn earworms and stink bugs apply 20 lbs. per acre. May be used for control of powdery mildew and rust. Start applications when insects appear or when diseases are reported in the area. Apply at 7-10 day intervals if needed. May be applied same day as harvest.

Do not graze treated fields or feed treated vines to dairy or meat animals. Sulphur may burn foliage when temperatures are excessively high.

### ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) .....	10%
Sulphur (as elemental) .....	50%
INERT INGREDIENTS .....	40%
TOTAL .....	100%

### CAUTION!

#### KEEP OUT OF REACH OF CHILDREN

(See additional Cautions and Directions)

HARMFUL IF INHALED OR SWALLOWED, DO NOT TAKE INTERNALLY. DO NOT BREATHE DUST. SKIN CONTACT MAY BE HARMFUL. AVOID CONTACT. WASH HANDS BEFORE EATING. TAKE SHOWER OR BATH AFTER WORK. WEAR REGULAR LONG SLEEVED CLOTHING. CHANGE TO CLEAN CLOTHING DAILY.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs and water receptacles.

For protection of honeybees avoid use if possible during periods when honeybees are visiting the crop or neighboring crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance (beyond bee flight range) until one week after application.

**COTTON—Early Season**—To control cotton fleahopper and thrips apply at the rate of 10 lbs per acre on appearance of insects.

**Mid and Late Season**—To control boll weevil, bollworm, cotton leafworms, lygus bugs, and cotton leaf perforator apply at the rate of 10 to 20 lbs. per acre per application. To control pink bollworm apply at the rate of 25 lbs. per acre per application. Apply at 5 to 7 day intervals on appearance of these insects.

**NOTE**—Aphid population will usually be repressed when this insecticide is used on a regular schedule of treatments.

Sevin does not control spider mites. Where infestations are encountered, use a recommended miticide.

Sevin may be applied after bolls open.

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

W. R. GRACE & CO.  
AGRICULTURAL CHEMICALS  
MEMPHIS, TENNESSEE

Amarillo Tex • Baltimore Md • Cary N.C. • Charleston S.C. • Columbus Ohio • Ft Pierce Fla • Henry Ill • John Mich • Lansing Mich  
Nashv Tenn • New Albany Ind • St Louis Mo

NET WT. 50 LBS.