

NACO
COPPER-SEVIN*-PARATHION 6-5-2 DUST

ACTIVE INGREDIENTS:

Copper as metallic (From Tribasic Copper Sulphate)	6.0%
Carbaryl (1-naphthyl N-methylcarbamate)	5.0%
Parathion (O,O-diethyl O-p-nitrophenyl thiophosphate)	2.0%
INERT INGREDIENTS	87.0%
TOTAL	100.0%

*Trademark of Union Carbide Corporation for 1-naphthyl N-methylcarbamate
U. S. Pat. Nos. 2,903,478; 3,009,855

(18-point) DANGER!
(12-point) KEEP OUT OF REACH OF CHILDREN
(See side panel for antidote and additional cautions and directions)




WARNING: POISONOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN! RAPIDLY ABSORBED THROUGH SKIN! DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

ANTIDOTES: If swallowed. Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet.

Call a Physician Immediately!

If on skin. In case of contact, remove contaminated clothing and immediately wash skin with soap and water.



This product is poisonous to fish and wildlife. Birds in treated areas will be harmed. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on this label. Dispose of containers and wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes. For protection of honeybees avoid use if possible during periods where honeybees are visiting the crop or neighboring crops. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance (beyond bee flight range) until one week after application.

DIRECTIONS FOR USE

Use the required amount of dust to cover foliage uniformly. Consult with your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with this label cautions and directions.

TOMATOES: For control of Aphids, fruitworms, hornworms, spider mites, early and late blight, and bacterial leaf spot apply at the rate of 25 - 35 lbs. per acre. Make the first application when the insects and/or disease appear and repeat at 7 - 10 day intervals as necessary.

Do not use within 10 days of harvest.



NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

W. R. GRACE & CO.
AGRICULTURAL PRODUCTS DIVISION
BALTIMORE, MARYLAND

NET WEIGHT POUNDS

ACCEPTED
May 14, 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND MITICIDICIDE ACT
FOR LISTING AND REGISTRATION
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.
2124-634