

NACO

NACO

USDA Reg. No. 2124-752

ACCEPTED
May 24-1971
2124-752



TOXAPHENE 6 E C

This product is poisonous to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, and ponds. Apply this product only as specified on this label. Repair equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

GENERAL DIRECTIONS

Naco Toxaphene 6 E C is especially formulated for the preparation of stable emulsions upon mixing with water. Water emulsions are easily applied by means of ground sprayers of high or low gallonage.

For ground sprayers apply not less than 5 gals. diluted spray per acre.

For airplane applications apply approximately 3 to 10 gallons per acre, consistent with crop growth and good coverage.

Consult with your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with this label cautions and directions.

COTTON: To control boll weevil apply 1 1/3 quarts per acre. Apply on foliage every 3 to 7 days until boll weevil is controlled. To control thrips apply 1 quart per acre. Apply on foliage as needed. Naco Toxaphene 6 E C will suppress cotton aphid when used throughout the season. Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.

ALFALFA: To control spittlebug nymphs use 1 quart per acre. Apply in early spring or after cutting before new growth is 4 inches tall.

CORN: To control armyworm use 1-1 1/3 quarts per acre. Apply when larvae are young.

To control grasshoppers use 1 quart per acre. Apply to hatching areas when nymphs are young.

Do not feed ensilage from treated corn to dairy animals or animals being finished for slaughter.

Do not graze dairy animals on treated stover.

ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene containing 67-69% Chlorine) 58.48%

Xylene 30.70%

INERT INGREDIENTS 10.82%

TOTAL 100.00%

WARNING!

KEEP OUT OF REACH OF CHILDREN

(See other warning statements on side panels)

MAY BE FATAL IF SWALLOWED. DO NOT BREATHE SPRAY MIST. DO NOT GET IN EYES, ON SKIN, OR ON CLOTHING. Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes get medical attention.

Do not store near feed or foodstuffs.

Not for use or storage in or around the home.

Keep away from heat and open flame.

DIRECTIONS (continued)

Do not graze meat animals on treated stover within 4 weeks of slaughter. No limitation on use of grain.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

SOYBEANS: To control velvetbean caterpillar, bean leaf beetle, armyworm and green cloverworm apply 3 pints of this product per acre. Do not apply within 21 days of harvest.

Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

SMALL GRAINS (Barley, Oats, Wheat and Rye): To control armyworm use 1 quart per acre. Apply when larvae are young.

To control grasshoppers use 1 quart per acre. Apply to hatching areas when nymphs are young.

Do not feed treated forage to dairy animals or animals being finished for slaughter. No restriction on the use of the grain.

TOMATOES: To control hornworms, tomato fruitworm, tomato pinworm, armyworm and climbing cutworm apply 3 to 6 pints of this product per acre.

Do not apply within 3 days of harvest.

ORNAMENTAL TREES AND SHRUBS: To control bagworm on arborvitae, juniper, spruce, pine, sycamore, willow and maple use 2 2/3 pints Toxaphene 6 E C per acre. Apply when bags are small.

Consult with your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with this label directions and cautions.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

W. R. GRACE & CO.
AGRICULTURAL CHEMICALS
MEMPHIS, TENNESSEE

Baltimore, Md. • Cary, N.C. • Columbus, Ohio • Charleston, S.C. • Fort Pierce, Fla. • Lansing, Mich. • New Albany, Ind. • Nashville, Tenn. • Ames, Iowa • Joplin, Mo. • Henry, Ill. • Amarillo, Tex.

NET CONTENTS GALLON(S)

