



NACO

EPA REG. NO. 2124-779

CAP-COP FUNGICIDE

ACTIVE INGREDIENTS:	BY WT.
Copper (in Basic Copper Sulfate) expressed as metallic	30.25%
*Captan	21.35%
INERT INGREDIENTS	48.40%
TOTAL	100.00%

*N- [(trichloromethyl) thio] - 4 cyclohexene-1, 2-dicarboximide
 U. S. Patent Nos. 2,553,770 - 2,553,771

Feb. 9, 1973

CAUTION!
KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. Avoid inhalation of dusts or spray mists. Avoid contact with skin. If taken internally, induce vomiting by giving mustard and water and tickling throat. Give large doses of magnesia or lime water, followed with doses of Epsom Salts.

Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers. Dispose of wastes and containers by burying in non-crop lands. Never re-use empty containers.

DIRECTIONS FOR USE

Naco Cap-Cop Fungicide may be used on peppers and tomatoes for control of the various diseases listed below. Use in conventional hydraulic sprayers mixing the required amount of this product in sufficient water for uniform coverage. Do not use in low-volume ground applicators. Do not combine with emulsifiable liquids unless previous use of the mixture has proven physically compatible and safe to plants. Consult with your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with this label cautions and directions.

Pour recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing and spraying operations. Do not allow mixture to stand.

PEPPERS: To control anthracnose, cercospora leaf spot (frog-eye spot and stem and rot) use 7 lbs. in 100 gals. water per acre. Apply at 3 to 7 day intervals throughout the growing season. To control early and late blight and damping off apply as a plant bed treatment. Begin when all seeds have germinated. Repeat at 10 day intervals.

TOMATOES: To control anthracnose, early and late blights, bacterial spot, gray leaf spot (stemphylium) and septoria leaf spot use 7 lbs. in 100 gals. water per acre. Begin in plant bed or field before diseases appear. Repeat at 4 to 14 day intervals.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

W. R. GRACE & CO.
AGRICULTURAL CHEMICALS
 MEMPHIS, TENN.

Baltimore, Md. • Cary, N.C. • Columbus, Ohio • Charleston, S.C. • Fort Pierce, Fla. • Lansing, Mich.
 New Albany, Ind. • Nashville, Tenn. • Ames, Iowa • Joplin, Mo. • Herrin, Ill. • Amarillo, Tex.

NET WEIGHT 7 LBS.