

(front panel)

GORDON'S

1% LINDANE DUST
FOR VEGETABLES

ACTIVE INGREDIENT:

Gamma Isomer of Benzene Hexachloride from Lindane.....	1%
<u>INERT INGREDIENTS.....</u>	<u>.99%</u>
TOTAL	100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panels for additional
precautionary statements and
Statement of Practical Treatment.

NET WEIGHT 50 POUNDS

ACCEPTED
with COMMENTS
in EPA Letter Dated:

MAR 2 1984

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

2217-179

STOP! READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLLOW
DIRECTIONS CAREFULLY.

NOTE: AERIAL APPLICATION OF LINDANE IS PROHIBITED.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals:

CAUTION: Harmful if swallowed. Avoid breathing dust. Avoid contact with eyes and skin. May be absorbed through skin. Wash thoroughly after handling. Avoid contamination of food and feedstuffs.

Statement of Practical Treatment:

Environmental Hazards:

This product is toxic to fish and wildlife. Do not apply directly to water. Do not apply when weather conditions favor drift from treatment area. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agriculture Extension Service.

Use Precautions:

This product should not be applied under conditions which might result in a toxic residue or residual odor or taste remaining on foodstuffs or edible portions of treated crops or forage at harvest time. Do not use on forage crops to be fed to dairy animals or to animals being finished for slaughtering.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Use 1% LINDANE DUST against the following pests on the listed plants. The usual rate of application is 25-30 lbs. per acre. However, when plants are small, as little as 10-15 lbs. per acre may be used and when plants are large, insects numerous, or when applied by airplane, up to 40 lbs. per acre may be required. Consult your State Agriculture Extension Service. Repeat applications as necessary. Note specific limitations under listed crops.

CELERY-Aphids, Thrips and Tarnished Plant Bugs. Apply before crop is half grown and before bunching.

COLE CROPS (CABBAGE, BROCCOLI)-Aphids, Thrips, Cutworms, Maggots. Apply only before heads or other edible parts begin to form.

CUCUMBERS, CANTALOUPE, WATERMELONS, SQUASH, PUMPKINS, MELONS-Cucumber Beetles, Aphids, Melon Worm, Leaf Miner, Thrips. Apply only before the blossoms open. Application to Melons, Pumpkins and Squash may result in reduction of quality at harvest time.

EGG PLANTS-Aphids, Leaf Miners, Lace Wing Bugs. Do not apply after edible parts start to form.

GLADIOLUS-Thrips.

ONIONS-Thrips. Make no applications to green or spring onions.

PEPPERS-Aphids, Leaf Miners. Apply only prior to blossom fall. Do not apply after edible parts start to form.

SPINACH-Aphids, Leaf Miners, Thrips, Loopers, Web Worms. Apply only to seedling Spinach.

STRAWBERRIES-Spittle Bugs, Strawberry Weevils. Apply only before blossoms open. Thoroughly treat the flower buds. Do not apply after fruit starts to form. Do not apply more than 15 pounds of 1% Lindane Dust per acre for foliage treatment.

TOMATOES-Colorado Potato Beetle, Flea Beetle, Aphids, Thrips. Apply only before blossoms fall. Do not apply after edible parts begin to form.

TOBACCO-Green Peach Aphids. Plant beds should be dusted at the rate of 20 lbs. per acre, repeating once a week.

STORAGE & DISPOSAL.

STORAGE: Store in original container in a locked storage area inaccessible to children and pets. To prevent cross-contamination during storage, keep separate from other pesticides, fertilizers, seeds, food, and feedstuffs.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY; DISCLAIMER. Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even where label directions are followed. Except where such disclaimers and exclusions are specifically prohibited by applicable law, MANUFACTURER MAKES NO WARRANTY OF MERCHANTABILITY, THERE ARE NO WARRANTIES WHICH EXTEND BEYOND THE DESCRIPTION ON THE FACE HEREOF, AND IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR ANY CONSEQUENTIAL DAMAGE OR LOSS.

EPA Reg. No. 2217-179

EPA Est. No. 2217-KS-1

Mfd. by  **pbi / GORDON**
CORPORATION
KANSAS CITY, KANSAS 66118

581

435/111912
37/1

M. J. ...

U.S. Environmental Protection Agency
Office of Pesticide Programs
Washington, D.C. 20460

Dear Mr. ...:

300/117581
17/2

Subject: Amendment - ...
Gordon's 1% Lindane Insecticide for Vegetables
EPA Registration No. 2217-173
Your Application dated November 15, 1977

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable, provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

1. Revise the statement of Practical Treatment to read as follows:

If swallowed - Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with a finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If inhaled - Remove victim to fresh air. Administer artificial respiration if indicated.

If on skin - Remove contaminated clothing and shoes. Wash areas with soap and water.

If in eyes - Flush eyes with plenty of water. Call a physician immediately.

2. Revise the label directions to read:

Do not apply to fields to be reseeded or to areas to be treated or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds growing in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

BEST DOCUMENT AVAILABLE

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							

2117-112

-2-

3. Delete "or when applied by airplane" from the general use directions paragraph.

Submit five (5) copies of your revised final printed labeling before you release the product for shipment with the amended labeling.

A stamped copy of the label is enclosed for your records.

If you hold a Federal registration, a supplemental registration, or a Special Local Needs registration, you may continue to produce new stocks of such products, using current labeling, until April 1, 1984. After that date, all remaining inventories of affected products will fall under the Existing Stocks provisions stated in our letter of October 28, 1983.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (78-767)

Enclosure

DCR-05425:LaRocca:2/2/84:RD-04:Rm. 217:557-2400:del. 2/19/84:pac

BEST DOCUMENT AVAILABLE