

December, 1966

NET WEIGHT 80 LBS.

(Kerr Logo)

10-10-10 + 0.20 D

CONTAINS 0.20% (R) DI-SYSTON OR 4 LBS. ACTIVE PER TON

For Use on Cabbage, Cauliflower, Broccoli & Brussel Sprouts

STATEMENT OF INGREDIENTS FOR INSECT CONTROL

ACTIVE INGREDIENTS:

O,O - Diethyl S- $\frac{1}{2}$ -(ethylthio) ethyl	
phosphorodithioate*	0.20%
INERT INGREDIENTS:	99.80%
TOTAL	100.00%

*U. S. Patent No. 2,759,010

GUARANTEED ANALYSIS FOR PLANT FOOD

Total Nitrogen (N)	10.00%
Available Phosphoric Acid	
(P ₂ O ₅)	10.00%
Soluble Potash (K ₂ O)	10.00%

DIRECTIONS FOR USE

DI-SYSTON is an effective systemic insecticide for early season crop protection which has been combined with fertilizer for efficient and uniform application. For best results apply at time of planting. Do not use on other crops used for food or forage. Use only according to label directions. Applications at rates higher than those shown on label may result in illegal crop residues.

VEGETABLES

CABBAGE AND BROCCOLI: For control of Aphids, Flea Beetles and Leafhoppers. Rate: Apply 500 lbs. (1 lb. of active insecticide) per acre on 36 in. row spacing. Apply as a band on each side of the seed furrow at planting or as a side-dressing after plants become established. Do not apply directly on the seed. Do not apply more than once per season. Do not apply within 14 days of harvest of Broccoli and within 42 days of harvest of Cabbage. Cabbage: For control of Root Aphids

Rate: Apply 750 lbs. (1- $\frac{1}{2}$ lbs. of active insecticide) per acre as outlined above.

BRUSSEL SPROUTS: For control of Aphids, Flea Beetles, Leafhoppers, Mites and Thrips.

Rate: Apply 500 lbs. (1 lb. of active insecticide) per acre on 36 in. row spacing.

Apply as a band on each side of the seed furrow at planting or as a side-dressing after plants become established. Do not

WARNING (18 Point Type)

KEEP OUT OF REACH OF CHILDREN

(12 Point Type)

Poisoning may occur as a result of exposure. May be fatal if swallowed. May be absorbed through skin. Do not breathe dust. Don't get in eyes, on skin, or clothing. Do not contaminate feed or food. Wash thoroughly after handling.

TO PHYSICIAN: Antidote: Atropine Sulfate. Atropine Sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

HIGHLY TOXIC TO FISH AND WILDLIFE. KEEP OUT OF LAKES, PONDS, AND STREAMS. DO NOT APPLY IN ANY MANNER NOT SPECIFIED ON LABEL.

DISPOSE OF WASTES BY BURYING IN NON-CROP LANDS AWAY FROM WATER SUPPLIES. CONTAINERS SHOULD BE DISPOSED BY BURNING (KEEP OUT OF SMOKE).

NOTICE

The Kerr-McGee Chemical Corp. and the seller make no warranty of any kind, express or implied, on the use of this material and assume no responsibility whatsoever for any personal injury, property damage, or other type of loss resulting from the use or handling of this material. The user assumes all risks for use or handling whether in accordance with directions or not.

The principle active ingredient of this product is patented under U. S. Patent No. 2,759,010 of Chemagro Corporation, Kansas City, Missouri, and the purchase of this product includes a limited license under said patent to use or reformulate the product only in the manner provided by the recommended

other crops used for food or forage. Use only according to label directions. Applications at rates higher than those shown on label may result in illegal crop residues.

VEGETABLES

CABBAGE AND BROCCOLI: For control of Aphids, Flea Beetles and Leafhoppers. Rate: Apply 500 lbs. (1 lb. of active insecticide) per acre on 36 in. row spacing. Apply as a band on each side of the seed furrow at planting or as a side-dressing after plants become established. Do not apply directly on the seed. Do not apply more than once per season. Do not apply within 14 days of harvest of Broccoli and within 42 days of harvest of Cabbage. Cabbage: For control of Root Aphids Rate: Apply 750 lbs. (1-½ lbs. of active insecticide) per acre as outlined above.

BRUSSEL SPROUTS: For control of Aphids, Flea Beetles, Leafhoppers, Mites and Thrips. Rate: Apply 500 lbs. (1 lb. of active insecticide) per acre on 36 in. row spacing. Apply as a band on each side of the seed furrow at planting or as a side-dressing after plants become established. Do not apply directly on the seed. Do not apply more than twice per season nor within 30 days of harvest. Allow a minimum of 21 days between applications.

CAULIFLOWER: For control of Aphids, Flea Beetles and Leafhoppers. Rate: Apply 500 lbs. (1 lb. active insecticide) per acre on 36 in. row spacing. Apply as a band on each side of the seed furrow at planting or as a side-dressing after plants become established. Do not apply directly on the seed. Do not apply more than twice per season. Do not apply within 40 days of harvest. Allow a minimum of 21 days between applications.

(R) DI-SYSTON is the Registered Trademark of Farbenfabriken Bayer A-G, Chemagro Corporation, licensee. U.S.D.A. Registration No. 9132-107

DISPOSE OF WASTES BY BURYING IN NON-CROP LANDS AWAY FROM WATER SUPPLIES. CONTAINERS SHOULD BE DISPOSED BY BURNING (KEEP OUT OF SMOKE).

NOTICE

The Kerr-McGee Chemical Corp. and the seller make no warranty of any kind, express or implied, on the use of this material and assume no responsibility whatsoever for any personal injury, property damage, or other type of loss resulting from the use or handling of this material. The user assumes all risks for use or handling whether in accordance with directions or not.

The principle active ingredient of this product is patented under U. S. Patent No. 2,759,010 of Chemagro Corporation, Kansas City, Missouri, and the purchase of this product includes a limited license under said patent to use or reformulate the product only in the manner provided by the recommended applications on the present label. The purchaser obligates itself to notify Chemagro Corporation of any proposed use or reformulation of such licensed product not provided by said recommended applications, and to obtain Chemagro's approval in writing before making any such use or reformulation of the licensed product. Any use, or reformulation intended for use of the licensed product in home and garden field is specifically prohibited by said license.

Manufactured By
KERR-McGee CHEMICAL CORP.
Oklahoma City, Okla.

