

3% THIODAN^R DUST

ACTIVE INGREDIENTS:

Endosulfan (Hexachlorohexahydromethano-
2,4,3-benzodioxathiepin oxide). . . 3.00%
INERT INGREDIENTS 97.00%
TOTAL 100.00%

(R) Trademark - Farbwerke Hoechst A.G.
U. S. Pat. No. 2,799,685

CAUTION (18 Pt. Type)

KEEP OUT OF REACH OF CHILDREN (12 Pt. Type)

Workers entering treated area on the day of application should wear protective clothing. Harmful if swallowed. Avoid inhalation or contact with skin. In case of contact with skin or eyes flush with plenty of water; for eyes get medical attention. Wash with soap and water after handling and before eating or smoking. Wear clean clothing. Do not apply or allow to drift to areas occupied by unprotected humans or domestic animals. Avoid contamination of feed and foodstuffs. THIS PRODUCT IS POISONOUS TO FISH AND WILDLIFE. KEEP OUT OF ANY BODY OF WATER. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT OF DUST FROM AREAS TREATED. CONTAINERS SHOULD BE DISPOSED BY PUNCHING HOLES IN THEM AND BURYING WITH WASTES, OR BY BURNING (KEEP OUT OF SMOKE).

DIRECTIONS FOR USE

BROCCOLI, CABBAGE, CAULIFLOWER: To control cabbage aphids, cabbage looper, imported cabbageworm, diamond back moth larvae, cross-striped cabbageworm. Flea beetle, harlequin bug and stink bug, use at the rate of 25-30 lbs. of dust per acre. Apply when insects or worms first appear and repeat as required. Do not apply to broccoli and cabbage within 7 days of harvest. Do not apply to cauliflower after the edible parts begin to form.

CUCUMBERS, PUMPKIN, MELONS, SUMMER AND WINTER SQUASH: To control aphids, cucumber beetles, striped cabbage flea beetle and squash vine borer, use at the rate of 25-30 lbs. per acre. Apply when insects first appear and repeat as required. For vine borer apply weekly to flower buds, stems and vines beginning when they first appear. May be applied 1 day before harvest.

DIRECTIONS FOR USE (Continued)

STRAWBERRIES: To control strawberry aphid and meadow spittle bug, use 20-30 lbs. per acre. Do not apply within 4 days of picking. Do not apply at intervals of less than 15 days, or more than twice during any 35 day period when fruit is present.

TOMATOES AND EGGPLANT: To control aphids, blister beetle, Colorado potato beetle, flea beetle, green stink bug, white fly and hornworm on tomatoes, use 20-30 lbs. per acre. Apply when insects first appear and repeat as needed. Do not apply within 1 day of picking.

CONDITIONS OF SALE

The seller guarantees the material sold to be true to the label. Further than that, it makes no representation or warranty, and it shall not be held responsible for any injury to crop, soil, person or property, resulting from the use or handling of said material, whether used in accordance with directions or not. The buyer and or user accept this material subject to the above conditions of sale.

Manufactured By
KERR-McGEE CHEMICAL CORP.
Oklahoma City, Okla.

U.S.D.A. Registration No. 9132-158

CAUTION (18 Pt. Type)

KEEP OUT OF REACH OF CHILDREN (12 Pt. Type)

Workers entering treated area on the day of application should wear protective clothing. Harmful if swallowed. Avoid inhalation or contact with skin. In case of contact with skin or eyes flush with plenty of water; for eyes get medical attention. Wash with soap and water after handling and before eating or smoking. Wear clean clothing. Do not apply or allow to drift to areas occupied by unprotected humans or domestic animals. Avoid contamination of feed and foodstuffs. THIS PRODUCT IS POISONOUS TO FISH AND WILDLIFE. KEEP OUT OF ANY BODY OF WATER. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT OF DUST FROM AREAS TREATED. CONTAINERS SHOULD BE DISPOSED BY PUNCHING HOLES IN THEM AND BURYING WITH WASTES, OR BY BURNING (KEEP OUT OF SMOKE).

DIRECTIONS FOR USE

BROCCOLI, CABBAGE, CAULIFLOWER: To control cabbage aphids, cabbage looper, imported cabbageworm, diamond back moth larvae, cross-striped cabbageworm. Flea beetle, harlequin bug and stink bug, use at the rate of 25-30 lbs. of dust per acre. Apply when insects or worms first appear and repeat as required. Do not apply to broccoli and cabbage within 7 days of harvest. Do not apply to cauliflower after the edible parts begin to form.

CUCUMBERS, PUMPKIN, MELONS, ^{squash} SUMMER AND WINTER SQUASH: To control aphids, cucumber beetles, striped cabbage flea beetle and squash vine borer, use at the rate of 25-30 lbs. per acre. Apply when insects first appear and repeat as required. For vine borer apply weekly to flower buds, stems and vines beginning when moths first appear. May be applied up to day of harvest.

PEPPERS: To control flea beetle and green peach aphid, use 30-35 lbs. per acre. Apply when insects first appear and repeat as required. Do not apply within 1 day of harvest.

POTATOES: To control Colorado potato beetle, flea beetle, leafhopper, Southern armyworm, European corn borer and three lined potato beetle, use 30-35 lbs. per acre. Apply as needed up to harvest.

DIRECTIONS FOR USE (Continued)

STRAWBERRIES: To control strawberry aphid and meadow spittle bug, use 20-30 lbs. per acre. Do not apply within 4 days of picking. Do not apply at intervals of less than 15 days, or more than twice during any 35 day period when fruit is present. **TOMATOES AND EGGPLANT:** To control aphids, blister beetle, Colorado potato beetle, flea beetle, green stink bug, white fly and hornworm on tomatoes, use 20-30 lbs. per acre. Apply when insects first appear and repeat as needed. Do not apply within 1 day of picking.

CONDITIONS OF SALE

The seller guarantees the material sold to be true to the label. Further than that, it makes no representation or warranty, and it shall not be held responsible for any injury to crop, soil, person or property, resulting from the use or handling of said material, whether used in accordance with directions or not. The buyer and or user accept this material subject to the above conditions of sale.

Manufactured By
KERR-McGEE CHEMICAL CORP.
Oklahoma City, Okla.

U.S.D.A. Registration No. 9132-158

