

USDA Reg. No. 2460-6

15% FERMATE* DUST

**28 LBS.
NET**

ACTIVE INGREDIENTS:
 Ferbam (Ferrie Dimethyl Dithiocarbamate)

INERT INGREDIENTS:
 Inerts in fermate
 Pyrophyllite clay
TOTAL

11.4%

2.7%
85.9%
100.0%

**FOR TOBACCO
PLANT BEDS**

DIRECTIONS— Start dusting when plants are dime size or when blue mold is reported in the area and dust twice a week until transplanted.

1. Apply early in the morning when the air is quiet and the plants moist with dew.
2. Do not fill the dust hepper more than 2/3 full at any time.
3. Be sure all leaves show a good coating of the dust.
4. In narrow beds (not over 3 yds. wide) with board sidewalls that hold the cover well above the plants, the first 3 or 4 applications of dust may be applied through the cover. For latter application and in other type beds always remove the cover before treating.

AMOUNT OF DUST REQUIRED TO COVER PLANTS:

1st to 4th applications	1/2 to 1 1/2 lbs. per 100 sq. yds.
5th and 6th applications	2 to 2 1/2 lbs. per 100 sq. yds.
All other applications	3 to 3 1/2 lbs. per 100 sq. yds.

ACCEPTED
 8-27-68
 2460-6

UNDER THE FEDERAL INSECTICIDE ACT
 OF 1947
 AS AMENDED
 BY PUBLIC LAW 85-623
 OCTOBER 3, 1958
 TO APPLY TO CERTAIN

CAUTION—May produce irritation of skin and mucous membrane. Avoid contact with eyes, skin or clothing. Avoid prolonged breathing of dust. In case of contact, flush immediately with plenty of water. For eyes get medical attention. **DANGER:** Keep away from fire and sparks. Avoid contamination of feed and feedstuffs.

"This product is toxic to fish. Keep out of lakes, streams, or ponds. Apply this product only as specified on this label."

* Trade Mark of DuPont Co.

CAUTION!

Keep Out of Reach
Of Children!

Howerton Gowen Chemicals, Inc.



NORFOLK, VA.

ROANOKE RAPIDS, N. C.

ROANOKE, VA.