

ACCEPTED

JUN 7 1990

MAX KILL

Contacta-Cide® 2

Odorless Contact Space Spray Stainless Quick Knockdown Ready To Use

PRECAUTIONARY STATEMENTS Hazards To Humans (and Domestic Animals)

CAUTION

Irritating to skin and eyes
Do not get in the eyes
Avoid skin contact
Wash thoroughly with soap and water after using
Avoid inhalation of spray mist

Do not remain in treated areas. Ventilate the area after treatment is completed.
Avoid contamination of cooking utensils, food ingredients, feed and water supplies.
Remove fish and birds and other domestic pets from room during spraying, and keep out of room until spray is dry.
Do not go by directly on pets, domestic livestock, especially dairy animals.
Do not give as internal treatment to animals.
Do not use in milk houses or dairy barns.
Do not apply directly to food.
Do not apply as a space spray while food processing is underway. Foods should be removed or covered before treatment. After spraying, wash all surfaces which will contact food with a suitable detergent and rinse with potable water.
With the exception of federally inspected meat and poultry plants, food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions indicated.
Destroy any contaminated food, feed or raw ingredients.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water. Do not contaminate water when disposing of equipment, wash waters. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Keep container closed when not in use.
Keep away from heat and open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MAX KILL® Contacta-Cide® 2 is a contact, READY-TO-USE insecticide. Use as directed on fish, birds, food mills, feed and food processing plants, saw-tooth grain processing plants, grain elevators, granaries, ship holds, barges, empty railcars, grain trucks, warehouses, feed and seed stores, poultry farms, restaurants and hotels.

KILLS THE ACCESSIBLE STAGES ON CONTACT:
Granary Weevil, Rice Weevil, Cadelle, Confused Flour Beetle, Saw-Toothed Grain Beetle, Angoumois Grain Moth, Indian Meal Moth, Cigarette Beetle, Drug Store Beetle, Lesser Grain Borer, Flies, Cockroaches, Silverfish, Spiders, Ants, Crickets, Gnats, Mosquitoes, *Hornets, *Wasps, *Mites.

ACTIVE INGREDIENTS
Pyrethrins 0.20%
*Piperonyl Butoxide, technical 0.40%
N-Octyl bicycloheptene dicarboximide . 0.66%

INERT INGREDIENTS 98.74%
*Equivalent to 0.32% of (Butylcarbityl) (6-Propylpiperonyl) ether and 0.08% of related compounds.

KEEP OUT OF THE REACH OF CHILDREN CAUTION
Statement of Practical Treatment

IF SWALLOWED: DO NOT INDUCE VOMITING. Seek medical aid. Aspiration of product into lungs due to vomiting can cause chemical pneumonia which can be fatal. Contains petroleum distillates.
IF BREATHED: If inhaled in heavy concentrations or if symptoms appear due to inhalation, remove to fresh air. Administer artificial respiration, oxygen or cardiopulmonary resuscitation if needed.
IF ON SKIN: Remove contaminated clothing. Wash with mild soap and water. Launder contaminated clothing. If irritation occurs, seek medical aid.
IF IN EYES: Flush with large amounts of water for at least 15 minutes. If irritation persists, seek medical attention.

See side panel for additional precautions.

NET CONTENTS GALLONS



MANUFACTURED BY
Research Products Company
Div. of McShores, Inc.
SALINA, KANSAS 67401

WHEN USED AS A SPRAY in mill, cereal grain processing plants, grain handling facilities (elevators, granaries) for the control of accessible stages of the granary weevil, rice weevil, cadelle, confused flour beetle, saw-toothed grain beetle, lesser grain borer and other listed crawling insects. Spray floors, walls and other surfaces of bins, storage and handling areas thoroughly with particular attention to cracks, crevices, and similar protected locations. Waste grain dust, dirt, and debris should be removed and destroyed and storage areas thoroughly cleaned before spraying. Treat unloading, handling and processing areas, as well as EMPTY conveying, processing and handling equipment for example the inside of reel & tiers, purifier conveyors, sifters, roll housings and top pans, pick-up conveyors, etc. Spray thoroughly around the base of heavy machinery and equipment. Let sprayed surfaces thoroughly dry before using. One gallon will cover 750 square feet of non-absorbent surface.

AS A PROTECTANT in empty boxcars, grain trucks, barges and shipholds for the control of accessible stages of the granary weevil, rice weevil, cadelle, confused flour beetle, saw-toothed grain beetle, lesser grain borer and other listed crawling insects. Spray the floors, walls, beams, cracks, crevices and other protected harborage points until surfaces are wet. A tanktype sprayer may be used utilizing low pressure. Allow for complete drying before loading. One gallon will cover 750 square feet of non-absorbent surface.

WHEN USED AS A SPRAY in food plants, grocery stores, hotels, feed and seed stores and restaurants for the control of accessible stages of cockroaches, silverfish, spiders, ants, crickets, and listed cereal grain insects, use a spray which can operate at a low pressure and which will produce a coarse non-misting spray pattern. Avoid contamination of food or foodstuffs as well as utensils or other food containers. Spray the floors, mold boards, cracks, crevices and other areas where insects are known to hide. (See other precautions on side panel.) Apply spray until areas are wet but not to point of runoff. One gallon will cover 750 square feet of non-absorbent surface.

WHEN USED AS A FOG for Indian meal moth, angoumois grain moth, hornets, wasps, gnats, mosquitoes, flies and accessible stages of lesser insects in mill, food processing plants, elevators and empty granaries or grain bins, use a mechanical or an operated fogging apparatus and MAX KILL® Contacta-Cide® 2 at the rate of one quart per 5,000 cubic feet. Close all doors and windows before application and leave closed for 30 minutes after total dosage has been applied to allow for settling.

WHEN USED AS A FOG in empty boxcars, van type grain trucks, barges and ship holds for the control of accessible stages of the granary weevil, rice weevil, cadelle, confused flour beetle, saw-toothed grain beetle, lesser grain borer and listed flying insects, use a mechanical or an operated fogging apparatus and MAX KILL® Contacta-Cide® 2 at the rate of one quart per 5,000 cubic feet. Close all doors and hatches before applying and leave closed for 30 minutes after total dosage has been applied to allow for settling.

WHEN USED AS A FOG in boxcars or van type trucks containing bagged flour or feed for the control of listed flying insects and accessible stages of listed cereal grain insects, cover the bags with rubber paper or plastic tarps and apply at the rate of 8 ounces (1/2 pint) per 1,000 cubic feet of unoccupied space. Close all doors before applying and leave closed for 30 minutes after total dosage has been applied.

STORAGE AND DISPOSAL

- 1. PROHIBITIONS**
Do not contaminate water, feed or food by storage or disposal.
- 2. PESTICIDE DISPOSAL**
Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
- 3. CONTAINER DISPOSAL**
Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.
- 4. GENERAL**
Consult federal, state and local disposal authorities for approved procedures other than those given above. Approved procedures vary considerably for different types of generators.
- 5. EQUIVALENT METHOD TO TRIPLE RINSING**
Empty drums as completely as possible. Remove the bungs and place the drums upside down until liquid has evaporated.

Non Warranty

Some make no warranty, expressed or implied, concerning the use of the product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.