

DM04 2724-487

9/14/98

1/3

**For Pest Control Operators and Commercial Use Only**

**ARTHITROL® 0.5% DURSBAN® PASTE BAIT WHICH KILLS  
ROACHES, ANTS, AND CRICKETS**

CONTAINS Bitrex®

**ACTIVE INGREDIENT:**

Chlorpyrifos (CAS #2921-88-2) ..... 0.5%

**OTHER INGREDIENTS:** ..... 99.5%

Total ..... 100.0%

**KEEP OUT OF REACH OF CHILDREN**

Do not apply to areas accessible to children or pets.

**CAUTION**

**PRECAUTIONARY STATEMENTS**


**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

May be harmful if swallowed. Do not take internally. Avoid eye contact. Wash thoroughly after handling. Wash contaminated clothing before reuse. Keep away from food, feedstuffs, and domestic water supplies.

See side panel for additional Precautionary Statements

Wellmark International  
Bensenville, IL 60106

EPA Reg. No. 2724-487



EPA Est. No. 2724-TX-1

**FIRST AID: IF SWALLOWED:** Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. **IF ON SKIN:** Wash thoroughly with soap and water. **IF IN EYES:** Flush with plenty of water. Get medical attention if irritation persists.

**NOTE TO PHYSICIAN:** Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine, by injection only, is antidotal.

**ENVIRONMENTAL HAZARDS:** This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

**STORAGE and DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. **Storage:** Store in original container out of reach of children and pets. Do not store above 100° F for extended periods of time. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of according to use directions or at an approved waste disposal facility.

**Container Disposal:** Wrap used container and place in trash can.

**DIRECTIONS FOR USE:** It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

ARTHITROL PASTE is a slow-drying insecticidal bait which is intended for the control of roaches, ants, and crickets.

MAY BE USED AS A SPOT OR CRACK AND CREVICE TREATMENT FOR THE INDOOR AND OUTDOOR CONTROL OF ANTS, COCKROACHES, AND CRICKETS IN AND AROUND DWELLINGS, BOTH HOMES AND APARTMENT BUILDINGS, AND INSTITUTIONAL, WAREHOUSE (INCLUDING WAREHOUSES THAT STORE CANNED, BOTTLED, OR PACKAGED FOOD ITEMS) AND COMMERCIAL FACILITIES INCLUDING OFFICE BUILDINGS, MANUFACTURING FACILITIES, HOTELS/MOTELS, RETAIL/WHOLESALE STORES (GROCERY STORES/SUPERMARKETS PET SHOPS), HOSPITALS, LABORATORIES, BUSES, BOATS/SHIPS, TRAINS, ZOOS, RESTAURANTS, AND FOOD PROCESSING PLANTS. **NOT FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY PLANTS.**

**INDOOR**

Indoor application sites are areas such as: entry areas, lavatories, locker rooms, restrooms, machine areas, boiler rooms, garages, closets, storage areas, and commercial and private food storage, preparation, and consumption areas, such as pantries, kitchens, dining rooms, and taverns/bars.

Keep exposed paste away from open food and food contact surfaces. Do not treat food preparation surfaces, or locations where routine operations, such as cleaning, may transfer diluted bait to food or to food preparation surfaces. In case of accidental exposure to food preparation surfaces, clean the surface with a detergent.

The recommended application rate for small cockroaches such as German or Brown banded is 1 to 2 pea-sized spots (each about 0.25 gm) of bait per 6 square feet. This rate should be doubled for large cockroaches such as the American cockroach. In heavy infestations, the rate should be increased to 3 to 12 inch beads of bait per 6 square feet. Bait must not be placed where it is accessible to animals or children.

Arthitrol Paste is packaged in a plastic syringe dispenser. For applications in cracks and crevices, apply by touching tip to surface to be treated, then depress plunger to dispense desired amount of product. Recap dispenser when application is complete.

Inspect intended application sites to determine locations frequented by pest(s). (Monitoring of the infestation using the Zoecon monitoring trap may be helpful.) Place bait where inspection indicates the pests frequent. Usual locations includespots, cracks, and crevices under and behind files, storage cabinets, and equipment; under refrigerators and sinks; in, under, and behind cabinets and tables; in closets; in void spaces; in attics; in electrical boxes (use caution when applying to electrical boxes); in pantries and food storage areas; and under water heaters. Numerous small placements of bait will provide better control than fewer larger placements. This is especially true for German cockroaches. In cases where heavy activity is indicated in void spaces that cannot be reached with Arthitrol Paste, Arthitrol Granular Bait may be used in the spaces that cannot be effectively reached with the paste.

Use caution when applying paste to porous surfaces such as wallpaper, wood, or paper, as possible staining may result. In questionable cases, place a small amount of bait in an inconspicuous location. Application of bait in areas of high temperature, such as near stoves and fryers, may cause bait to run.

Foraging trails are effective baiting sites for ants. When baiting for ants, apply around trails and foraging areas. Foraging workers will transport the product into the colony and down to the brood and queen(s). Colony kill times may vary from a few days to six weeks. Retreatment is desirable for large mounds and in the event of rains.

## **OUTDOOR**

For outdoor baiting, use recommended indoor rates and place in areas adjacent to the structure, at potential access points such as foundation cracks, soffits, sills, pipe and wire penetrations, expansion joints and eaves, in/around wood piles, around windows, doors, sashes, storage buildings, siding, shingles, eaves, weep holes, and water meter boxes. Concentrate applications at point of entry areas.

For heavy infestations, a reinspection of the treated area is recommended between one week and one month after initial bait placement. Replace bait as necessary if visibly depleted and the pest(s) are still present. You should notice visible reduction in the pest(s) within a few days after application.

For information or in case of an emergency, call [phone number].

Seller makes no warranty, express or implied, concerning the use and handling of this product other than indicated on the label. Buyer assumes all risks of use and handling of this material when such use and handling are not in accordance with directions.