

P21  
PUE

Front Panel

**ACCEPTED**  
JUN 8 1990  
U.S. DEPARTMENT OF AGRICULTURE  
Federal Insecticide, Fungicide, and  
Rodenticide Act, as amended, for the  
purpose mentioned on the  
label.

2900-18

**SHEPARD'S  
LIQUID WATER BUG AND FLEA KILLER**

A Ready-to Use Insecticide with Flushing and  
Quick Knockdown Characteristics

2900-18

**KILLS WATERBUGS, ROACHES, FLEAS AND  
BROWN DOG TICKS**

For Use in Homes, Garages and the Non-Food Areas of  
Motels, Hotels, Institutions, Warehouses

**Active Ingredients:**

Pyrethrins .....	0.052%
*Piperonyl butoxide, Technical .....	0.260%
**Diazinon .....	0.500%
Inert Ingredients: .....	99.188%
Total 100.000%	

\* Equivalent to min. 0.208% (butylcarbityl) (6-propylpiperonyl) ether and 0.052% related compounds.

\*\* 0,0 diethyl 0-(2-isopropyl-6-methyl-4 pyrimidinyl) phosphorothioate.

Keep Out of Reach of Children

**CAUTION**

See Side Panel for Additional Precautionary Statements

EPA Reg. No. 2900-18

EPA Est. No. 2900-NC-1

Do Not Transport or Store Under 32° F.

Net Contents: 1 Gallon

Sold by  
SHEPARD CHEMICAL WORKS  
PO Box 473  
Wilmington, NC 28402

**BEST AVAILABLE COPY**

Left Panel

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND  
DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. May be absorbed through the skin. Avoid breathing of spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling and before smoking or eating. Do not allow spray to contact food, feedstuffs or water supplies. Thoroughly wash dishes and food handling utensils if contaminated by this spray. Do not use on household pets or humans.

Do not spray in the immediate area when others are present. After application, open windows and/or doors to provide adequate fresh air ventilation to the room(s). Remove clothing after spraying and launder before reuse.

Do not allow children or pets into a treated area until the spray has dried.

Statement of Practical Treatment

If Swallowed: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If Inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.

If on Skin: Remove contaminated clothing and wash affected areas with soap and water.

If in Eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Note to Physician: If symptoms of cholinesterase inhibition are present, atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**BEST AVAILABLE COPY**

## Right Panel

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### To Kill Waterbugs and Roaches, Ants and Spiders:

Spray areas where these pests are found or may hide, including dark corners of rooms, closets, cracks, crevices, along baseboards, and door sills and frames, behind and beneath sinks, stoves, refrigerators, cabinets and around garbage cans, plumbing and other utility installations. Pests flushed out of hiding should be sprayed directly. Treat points of entry such as around cellar window sills and frames, door sills and frames and between foundations and house sills.

Repeat treatment as necessary.

#### To Kill Fleas and Brown Dog Ticks, Indoors:

Clean premises thoroughly before treatment. Using a coarse cone or fan spray, treat the entire floor and carpet area. Special attention should be given to cracks, crevices and under carpet and rug edges. Also treat animal sleeping quarters, closets, upholstered furniture (with particular attention to seams), beds, under sofas and other furniture. In animal sleeping quarters, bedding should be cleaned or replaced with fresh bedding. Do not use this application in nurseries or any area where infants may sleep. Apply at the rate of approximately 1 gallon to 750 square feet (17 fl. oz. per 100 sq. ft.). To avoid reinfestation, pets must be treated simultaneously with an EPA approved insecticide labeled for use on animals. Application should be repeated as necessary. Humans and pets should avoid contact with treated surfaces (rugs, upholstered furniture, etc.) for 4 hours after treatment.

Do not use in food areas of food handling establishments, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Non-food areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, garages, mop closets and storage (after canning or bottling).

Not for Use in USDA Meat and Poultry Plants.

**BEST AVAILABLE COPY**

02-10-7

Right Panel (cont)

STORAGE AND DISPOSAL

Securely wrap original container in several layers of newspaper and discard in trash. Do not reuse container. Rinse thoroughly before discarding in trash.

Storage: Store in original container and place in locked storage area.

WORK SAFETY RULES

Repeated exposures to cholinesterase inhibitors such as are contained in this product may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor.

If handled indoors, provide mechanical exhaust ventilation.

Keep all unprotected persons, children, livestock, and pets away from treated areas or where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way, stop work and get help right away. See Practical Treatment Section.

**BEST AVAILABLE COPY**