

DIRECTIONS

COTTON: Boll Weevils, on small cotton apply 1 quart Toxaphene Emulsion per acre diluted with enough water to give sufficient coverage. On large cotton apply 2 quarts Toxaphene Emulsion per acre in sufficient water to give good coverage. On large cotton at least 3 spray nozzles per row is recommended. Nozzles should be kept at least 10 inches from the plants to avoid leaf burn. Begin applications at 5-day intervals when (1) 1 or more adult weevils are found per 100 terminals 1 week before squaring (8-leaf stage), or (2) the square infestation first rises to 10% (if pre-square applications were not needed). Continue applications as long as the plants are squaring. For best results, keep spray mixture agitated. For safety, the spray boom on ground equipment should be located behind the operator.

THIS PRODUCT IS NOT TO BE USED FOR USE OF...

CARTER
Carolina
PEANUT

Toxaphene

EMULSION (UNREFINED)

ACTIVE INGREDIENTS:

Toxaphene*	60.0%
INERT INGREDIENTS	40.0%
*Technical Chlorinated Camphene-Chlorine 67% to 69% TOTAL -- 100.0%	
Each Gallon contains 6 lbs. Technical Toxaphene	

1 GALLON NET

Packed & Distributed By

CARTER INSECTICIDE & CHEMICAL CO., INC.

P. O. Box 209 Wallace, N. C. 28166 Phone AT 5-2111

WARNING

NOTICE: Conditions of Sale

Small, faint text detailing the conditions of sale for the product.

Handwritten date: 5/1/57


RECEIVED WITH COMMENTS

DIRECTIONS

COTTON: Boll Weevils, on small cotton apply 1 quart Toxaphene Emulsion per

sufficient coverage. On large cotton apply 2 quarts Toxaphene Emulsion per acre in sufficient water to give good coverage. On large cotton at least 3 spray nozzles per row is recommended. Nozzles should be kept at least 10 inches from the plants to avoid leaf burn. Begin applications at 5-day intervals when (1) 1 or more adult weevils are found per 100 terminals 1 week before squaring (8-leaf stage), or (2) the square infestation first rises to 10% (if pre-square applications were not needed). Continue applications as long as the plants are squaring. For best results, keep spray mixture agitated. For safety, the spray boom on ground equipment should be located behind the operator.

THIS PRODUCT IS NOT RECOMMENDED FOR USE ON TOBACCO.



Toxaphene

EMULSION CONCENTRATE

ACTIVE INGREDIENTS:

Toxaphene*	60.0%
INERT INGREDIENTS	40.0%
*Technical Chlorinated Camphene-Chlorine 67% to 69% TOTAL — 100.0%	
Each Gallon contains 6 lbs. Technical Toxaphene.	

WARNING

KEEP OUT OF REACH OF CHILDREN & ANIMALS
See right panel for additional warnings

1 GALLON NET

Packed & Distributed By
CARTER INSECTICIDE & CHEMICAL CO., INC.
P. O. Box 209 Wallace, N. C. 28466 Phone AT 5-2144

WARNING

May Be Fatal If Swallowed! Do not breathe spray. Do not get on face or on clothing. Do not graze or feed cotton trash to dairy animals or animals being finished for slaughter. Apply this product only as specified on this label.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning (keep out of smoke).

NOTICE: Conditions of Sale

Manufacturer Seller warrants that, unless otherwise specified herein, the products sold hereunder shall meet Manufacturer Seller's standard specifications and shall be of merchantable quality. Except as otherwise expressly stated herein, Manufacturer/Seller makes no representation or warranty of any kind, express or implied, with respect to the product. All claims for alleged defective products, shortage, or other cause shall be deemed waived unless made in writing and received by Manufacturer Seller within 20 days after Buyer's receipt of the product.

E-1137-A

USDA REG. NO. 2917-43

ACCEPTED
5/1/67
CARTER INSECTICIDE AND CHEMICAL CO.

ACCEPTED WITH COMMENTS