


Reg # 2935-420

PM-25

195



NUSAN FLOWABLE 30

ACTIVE INGREDIENT:	
2-(thiocyanomethylthio)benzothiazole.....	30
INERT INGREDIENTS:.....	70
TOTAL.....	100

EPA REG. NO. 2935-420

KEEP OUT OF REACH OF CHILDREN
WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash promptly and thoroughly with an abrasive soap and cool water and finally with glycerin. In case of persistent irritation of the skin, obtain medical attention.

IF IN EYES: Flood eyes immediately and thoroughly with cool water for 15 to 30 minutes. See a physician if any irritation persists.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

WARNING
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes eye damage. May be harmful if absorbed through the skin. Do not get in eyes, on skin or on clothing. Wear goggles or a face shield and rubber gloves when handling. Harmful if swallowed. Avoid contamination of food.

IMPORTANT! Do not use crops grown from NUSAN FLOWABLE 30 treated seed for grazing of livestock or to harvest as silage or forage for seven weeks after planting. However, grain harvested from plants grown from NUSAN FLOWABLE 30 treated seed can be used for food or feed.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

BEST COPY AVAILABLE

195

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water. Do not contaminate water by cleaning of equipment or disposal of wastes. For terrestrial uses, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark.

GENERAL INFORMATION

NUSAN FLOWABLE 30 is recommended for the control of soil- and seed-borne fungi which impair the germination and seedling development of FIELD CORN, OATS, RICE, SAFFLOWER, SORGHUM or MILO, SUGAR BEETS and WHEAT. NUSAN FLOWABLE 30, when used as directed, will usually increase stands and yields of field corn, oats, rice, sorghum or milo, sugar beets, safflower and wheat by preventing seed decay and seedling diseases.

NUSAN FLOWABLE 30 will also be used for the control of covered smut of barley and Helminthosporium stripe of barley; loose and covered smut of oats; covered kernel smut of sorghum; and stinking smut or bunt of wheat.

NUSAN FLOWABLE 30 is used at time of planting to protect cottonseed and seedlings against seed and seedling rots, damping off, soreshin (Rhizoctonia), Pythium, and other diseases impairing good seed germination and seedling development.

NUSAN FLOWABLE 30 is used at time of planting to protect safflower seed and seedlings against safflower rust (Puccinia carthami). NUSAN FLOWABLE 30 is also used to protect sugar beet seed and seedlings against seed and seedling rots (Phome batae) and damping-off (Pythium and Rhizoctonia). Use only higher rates under conditions of high pest pressure. Lower rates may give reduced efficiency.

Seed should be planted within 5 months of treatment. If stored longer than 5 months, re-treatment with NUSAN FLOWABLE 30 may be necessary.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on agricultural establishments in hopper-box, planter box, slurry box or other seed treatment application at or immediately before planting.

ALWAYS READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

The purchaser of this product is responsible for insuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21 CFR Part 2.25. Any dye added must be cleared for use under 40 CFR Part 180.1001.

Label treated seed. Seeds treated with NUSAN FLOWABLE 30 cannot be used for food, feed, oil production, or any other purpose except planting.

RECOMMENDATIONS FOR USE

BARLEY, OATS, RICE and WHEAT: Seed protection - For ready-mix treaters, use 0.75 fluid ounces of NUSAN FLOWABLE 30 per bushel of seed. NUSAN FLOWABLE 30 may be dispersed in water for distribution, provided the recommended amount of product is applied and the dispersion is agitated. For slurry treaters, disperse 0.75 fluid ounces of NUSAN FLOWABLE 30 in 4 fluid ounces of water and apply this dispersion per bushel of seed. However, the concentration and amount of slurry to be used must be adjusted to the specific conditions of the slurry treater for optimum seed coating, provided that 0.75 fluid ounces of NUSAN FLOWABLE 30 is applied per bushel of seed.

FIELD CORN: Seed protection - For ready-mix treaters, use 2.0 fluid ounces of NUSAN FLOWABLE 30 per 100 pounds of corn seed. NUSAN FLOWABLE 30 may be dispersed in water for distribution, provided the recommended amount of product is applied and the dispersion is agitated. For slurry treaters, disperse 2.0 fluid ounces of NUSAN FLOWABLE 30 in 10 fluid ounces of water and apply this dispersion per 100 pounds of seed. However, the concentration and amount of slurry to be used must be adjusted to the specific conditions of the slurry treater for optimum seed coating, provided that 2.0 fluid ounces of NUSAN FLOWABLE 30 is applied per 100 pounds of seed.

COTTON*: Seed protection - For ready-mix treaters, use NUSAN FLOWABLE 30 undiluted at the rate of 4.0 to 5.0 fluid ounces per 100 pounds of acid-delinted seed and 5.0 to 6.0 fluid ounces per 100 pounds of machine-delinted seed. NUSAN FLOWABLE 30 may be diluted with water for distribution, provided the recommended amount of product is applied. For slurry treaters, prepare slurry by stirring 1.6 to 2.0 pints (for acid-delinted seed) or 2.0 to 2.4 pints (for machine-delinted seed) of NUSAN FLOWABLE 30 in water to a final volume of 1 gallon. Suggested use is 0.125 pint of slurry per 10 pounds of seed. However, the concentration and amount of slurry to use must be adjusted to the specific conditions of the slurry treater provided that NUSAN FLOWABLE 30 is applied at 4.0 to 5.0 fluid ounces per 100 pounds of acid-delinted seed or 5.0 to 6.0 fluid ounces of machine-delinted seed. Texas only : use 1.25 to 6.0 fluid ounces per 100 pounds of seed.

*Under conditions of heavy pythium pressure and rates below 4

fluid ounces per 100 pounds of seed, WILBUR-ELLIS Company recommends treatment with WILBUR-ELLIS Apron TL or WILBUR-ELLIS Apron Flowable for additional systemic pythium control.

SAFFLOWER: Seed protection - For ready-mix treaters, use NUSAN FLOWABLE 30 at the rate of 1.0 to 3.0 fluid ounces per 100 pounds of seed. NUSAN FLOWABLE 30 may be dispersed in water for distribution, provided the recommended amount of product is applied and the dispersion is agitated. For slurry treaters, disperse 1.0 to 3.0 fluid ounces of NUSAN FLOWABLE 30 in 8 fluid ounces of water and apply this dispersion per 100 pounds of seed. However, the concentration and amount of slurry to use must be adjusted to the specific conditions of the slurry treater for optimum seed coating, provided that NUSAN FLOWABLE 30 is applied at 1.0 to 3.0 fluid ounces per 100 pounds of seed.

SORGHUM (Milo): Seed protection - For ready-mix treaters, use NUSAN FLOWABLE 30 at the rate of 1.25 to 2.5 fluid ounces per 100 pounds of seed. NUSAN FLOWABLE 30 may be dispersed in water for distribution, provided the recommended amount of product is applied and the dispersion is agitated. For slurry treaters, disperse 1.25 to 2.5 fluid ounces of NUSAN FLOWABLE 30 in 8 fluid ounces of water and apply this dispersion per 100 pounds of seed. However, the concentration and amount of slurry must be adjusted to the specific conditions of the slurry treater for optimum seed coating, provided that 1.25 to 2.5 fluid ounces of NUSAN FLOWABLE 30 is applied per 100 pounds of seed. California: Use only higher rate (2.5 fluid ounces).

SUGAR BEETS: Seed protection - For ready-mix treaters, use NUSAN FLOWABLE 30 at the rate of 3.0 to 4.0 fluid ounces per 100 pounds of seed. NUSAN FLOWABLE 30 may be dispersed in water for distribution, provided the recommended amount of product is applied and the dispersion is agitated. For slurry treaters, disperse 3.0 to 4.0 fluid ounces of NUSAN FLOWABLE 30 in 8 fluid ounces of water and apply this dispersion per 100 pounds of seed. However, the concentration and amount of slurry to use must be adjusted to the specific conditions of the slurry treater for optimum seed coating, provided that NUSAN FLOWABLE 30 is applied at 3.0 to 4.0 fluid ounces per 100 pounds of seed.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

595

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

STORAGE: Store in original container only and keep sealed. Store in closed storage areas. Use caution when moving, opening, closing or pouring.

NOTICE - READ CAREFULLY
CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

THE STATEMENTS MADE ON THIS LABEL ARE BELIEVED TO BE TRUE AND ACCURATE, BUT BECAUSE OF CONDITIONS OF USE WHICH ARE BEYOND OUR CONTROL, WILBUR-ELLIS COMPANY DOES NOT MAKE, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY WARRANTY, GUARANTY OR REPRESENTATION, EXPRESSED OR IMPLIED, CONCERNING THIS MATERIAL OR THE USE THEREOF, EXCEPT IN CONFORMITY WITH THE STATEMENTS ON THE LABEL. NEITHER WILBUR-ELLIS COMPANY NOR THE SELLER SHALL BE HELD RESPONSIBLE IN ANY MANNER FOR ANY PERSONAL INJURY OR PROPERTY DAMAGE OR LOSS RESULTING TO THE BUYER OR TO THE OTHER PERSON FROM HANDLING, STORAGE OR USE OF THIS MATERIAL, NOT IN ACCORDANCE WITH DIRECTIONS. THE BUYER ASSUMES ALL RISK AND LIABILITY RESULTING FROM IMPROPER HANDLING, STORAGE OR USE AND ACCEPTS AND USES THIS MATERIAL ON THESE CONDITIONS.

TREATMENT OF HIGHLY MECHANICALLY DAMAGED SEED OR SEED OF KNOWN LOW VIGOR AND POOR QUALITY, EXCEPT FOR THE PURPOSES OF CURATIVE CONTROL OF EXISTING DISEASE OR INSECT PESTS MAY RESULT IN REDUCED GERMINATION AND/OR REDUCED VIGOR. TREAT AND CONDUCT GERMINATION TESTS ON A SMALL PORTION OF SEED BEFORE COMMITTING THE TOTAL SEED LOT TO A SELECTED CHEMICAL TREATMENT. DUE TO SEED QUALITY CONDITIONS BEYOND THE CONTROL OF WILBUR-ELLIS COMPANY, NO CLAIMS ARE MADE TO GUARANTEE GERMINATION OF CARRY-OVER SEED.

WILBUR-ELLIS logo^R, and IDEAS TO GROW WITH^R are registered trademarks of WILBUR-ELLIS Company.

IN CASE OF EMERGENCY, CALL CHEMTREC: (800)424-9300

Manufactured By:
WILBUR-ELLIS COMPANY
Fresno, California 93755

BEST COPY AVAILABLE

5