

### **Directions For Use**

For the control of accessible stages of grain weevils, flour beetles and moth larvae, cadelle, saw toothed grain beetle, roaches, silverfish, ants, spiders, carpet beetles, cheese and flour mites, flies and flying moths.

As A Contact Spray. For the control of crawling insects listed above, apply directly onto walls, partitions, floors, in cracks, crevices, under skids, equipment, scales, motor mounts, on ledges, in conveyor, proofer, behind switch boxes, tanks, storage cabinets, floor drains, locker and lunch rooms and all places where insects may harbor and multiply.

Clean areas thoroughly before spraying. Dust and dirt absorb insecticide which limits effectiveness. Repeat treatment when necessary.

*P-B*

# **INSECT SPRAY**

## **FOR INSECT CONTROL IN FOOD PLANTS**

### **ACTIVE INGREDIENTS**

Pyrethrins	0.25%
Technical piperonyl butoxide*	0.20%
Petroleum Solvent	99.55
	TOTAL 100.00%

\*Equivalent to 16.00% of (butylcarboxylic 16% pyrethrins) and 0.80% of related compounds.

USDA REC No. 2993-56

### **— WARNING —**


#### **KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed. Do not spray near or toward open flame. Do not pour, spill or store near fire or flame. Avoid contamination of food and food processing equipment. Do not use inside Food Plant while Plants are in operation.

Seller makes no warranty expressed or implied, concerning the use of the product other than indicated on the label. Buyer assumes all risk of use and/or handling of the material when such use and/or handling is contrary to label instruction.

FLUID CONTENTS

GALLONS



### **Directions For Use**

As A Contact Spray. For the control of crawling insects listed above, apply directly onto walls, partitions, floors, in cracks, crevices, under skids, equipment, scales, motor mounts, on ledges, in conveyor, proofer, behind switch boxes, tanks, storage cabinets, floor drains, locker and lunch rooms and all places where insects may harbor and multiply.

Clean areas thoroughly before spraying. Dust and dirt absorb insecticide which limits effectiveness. Repeat treatment when necessary.

Do not allow residues, dust and sprayer during cleaning operations. Keep closed for 10-15 minutes after application. Do not treat and clean areas simultaneously.

Do not use inside Food Plant while Plants are in operation.

Do not use inside Food Plant while Plants are in operation.

**Brayton Chemicals, Inc.**  
BURLINGTON, IOWA

**Brayton Chemicals, Inc.**  
BURLINGTON, IOWA

**Brayton Chemicals, Inc.**  
BURLINGTON, IOWA