

KILCOP "53" WETTABLE

53% Copper

ACTIVE INGREDIENTS:

Copper as metallic, not less than 53.00%
(From basic copper sulphate)

INERT INGREDIENTS:

Not more than 47.00%

A NEUTRAL COPPER FUNGICIDE FOR FLORIDA

CAUTION

KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED

NON-WARRANTY: The Manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

USDA Reg. No. 3267 - 1

The Kilgore Seed Company

Division of Agrow Seed Company
PLANT CITY, FLORIDA

Reg. No. LC 011265



KILCOP "53"

SPRAYING RECOMMENDATIONS

The following recommendations for KILCOP "53" are based on pounds in 100 gallons of water as sprays. Apply spray to vegetable crops at a rate of 100 to 150 gallons per acre if conventional type spray equipment is used. Frequent applications throughout the growing season are usually required. However since the number of applications and time intervals depend on crop, disease, growth conditions and weather, ask Federal or State authorities for recommendations best suited to local conditions.

CELERY - EARLY BLIGHT: As a spray use 4 lbs. Begin applications when plants are established. Apply at weekly intervals.
EGGPLANT - PHOMOPSIS BLIGHT: As a spray use 4 lbs. Begin applications when plants are established. Apply at 7-10 day intervals.

PEPPERS - CERCOSPORA and BACTERIAL LEAF SPOTS: As a spray use 4 lbs. Begin spraying when first true leaves appear. Apply at weekly intervals or oftener during rainy weather.

POTATOES - EARLY AND LATE BLIGHTS: As a spray use 4 lbs. Apply every 7 to 10 days from time plants come up until maturing.

STRAWBERRIES - LEAF SPOTS: As a spray use 3 lbs. Usually 2 or 3 applications at 7 day intervals when leaf spots appear.
TOMATOES - EARLY and LATE BLIGHT; BACTERIAL LEAF SPOT: As a spray use 4 lbs. Begin applications when first true leaves appear and apply at weekly intervals until maturity.

WATERMELONS - ANTHRACNOSE: As a spray use 2 lbs. Begin applications when plants are established. Apply at 7 to 10 day intervals.

CITRUS - GREASY SPOT (Florida only): As a spray use ¾ lb. Apply between July 1 and August 10.

MELANOSE: Use 1½ lbs. A single spray applied 1 to 3 weeks after petal fall.

SCAB: Use 1½ lbs. Make 2 applications, the first applied during the dormant period and the second after 2/3 of the petals have fallen. If scab is severe, use 2 to 3 lbs. in the dormant spray.



ACCEPTED

APR 14 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 3267-1