

# 10% SEVIN DUST

**ACCEPTED**  
 4-26-71  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC POISON REGISTERED  
 UNDER NO. 3342-69

*Comments*

**ACTIVE INGREDIENTS:**

Carbaryl (1-naphthyl N-methylcarbamate).....10.00%

**INERT INGREDIENTS** .....90.00%

**TOTAL 100.00%**

**CAUTION: READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.**

**DIRECTIONS FOR USE**

**TOBACCO:** For the control of Flea beetles, Japanese beetles, Hornworm use 15 lbs. per acre. Grasshoppers 15 lbs. per acre. Apply dust evenly to tobacco foliage and to barrier strips 15 to 30 ft. wide around tobacco. Do not graze or feed animals on treated barrier strips for 7 days following treatment. Good coverage on underside of leaves over entire plant especially lower leaves is essential. Larger worms cause more damage and are harder to kill than small worms. Treat infested fields before worms exceed 1 1/2 inches in length.

**NOTE:** If late applications are necessary, treat immediately after priming rather than before priming. Treated fields may be entered immediately after dust has settled. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

**BEANS:** (Green beans, lima beans, snap beans, cowpeas and black eyed peas). Use 5 lbs. per acre for Mexican bean Beetle; use 10 lbs. per acre for bean leaf beetle, cucumber beetles, flea beetle, Japanese beetle, leafhoppers, velvet bean caterpillar and western bean cutworm, use 10 to 15 lbs. per acre for armyworms, corn earworm, stink bugs and tarnished plant bug. Use up to day of harvest. **CORN:** Use 10 to 20 lbs. per acre for corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetle, sap beetles and leafhoppers. For larvae in whorl (budworm damage) & foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. Use up to the day of harvest.

**CAUTION:** "Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations."

**CABBAGE, BROCCOLI, HEAD LETTUCE, CARROTS, POTATO, TOMATO, EGGPLANT:** Use 5 to 10 lbs. per acre for flea beetles and harlequin bug. 10 to 15 lbs. per acre for cabbage caterpillars, armyworms, & corn earworms. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity.

**CAUTION:** Do not apply within three days of harvest on Cabbage, Head Lettuce and Broccoli. Cucumber beetles, thrips, cutworms, leafhoppers, squash bugs, and squash borer on cucumbers, melons, pumpkins and squash apply 10 lbs. per acre."

**Pepper:** Flea beetles 10 to 15 lbs. per acre.

**Okra:** For the control of Blister beetles and Japanese beetles use 20 lbs. per acre.

**NOTICE:** Our recommendation for use of this product are based upon test believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such or in combination with other materials.

**CAUTION:** This product is highly toxic to fish and aquatic life. Do not apply to water bodies. Do not apply to areas where it may be carried into water bodies. Do not apply to areas where it may be carried into water bodies. Do not apply to areas where it may be carried into water bodies.

## CAUTION: KEEP OUT OF REACH OF CHILDREN AND ANIMALS

**CAUTION:** This product is highly toxic to fish and aquatic life. Do not apply to water bodies. Do not apply to areas where it may be carried into water bodies. Do not apply to areas where it may be carried into water bodies. Do not apply to areas where it may be carried into water bodies.

**CONTAINER DISPOSAL**

Destroy this container by cutting it lengthwise and burning it with the waste in burning. Do not reuse. Do not use for any other purpose. Do not use for any other purpose. Do not use for any other purpose.

**ACCEPTED**  
 4-26-71  
 UNDER THE FEDERAL INSECTICIDE ACT  
 FOR ECONOMIC ACTION REGISTERED  
 ETO NUMBER NO 3342-69

*Lim... ..*

**ACTIVE INGREDIENTS:**

Carbaryl (1-naphthyl N-methylcarbamate) .....

10.00%

**INERT INGREDIENTS** .....

90.00%

**TOTAL 100.00%**

**DIRECTIONS FOR USE**

**TOBACCO:** For the control of Flea beetles, Japanese beetles, Hornworm use 15 lbs. per acre. Grasshoppers 15 lbs. per acre. Apply dust evenly to tobacco foliage and to barrier strips 15 to 30 ft. wide around tobacco. Do not graze or feed animals on treated barrier strips for 7 days following treatment. Good coverage on underside of leaves over entire plant especially lower leaves is essential. Larger worms cause more damage and are harder to kill than small worms. Treat infested fields before worms exceed 1 1/2 inches in length.

**NOTE:** If late applications are necessary, treat immediately after priming rather than before priming. Treated fields may be entered immediately after dust has settled. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

**BEANS:** (Green beans, Lima beans, snap beans, cowpeas and black-eyed peas). Use 5 lbs. per acre for Mexican bean Beetle, use 10 lbs. per acre for bean leaf beetle, cucumber beetles, flea beetle, Japanese beetle, leafhoppers, velvet bean caterpillar and western bean cutworm, use 10 to 15 lbs. per acre for armyworms, corn earworm, stink bugs and tarnished plant bug. Use up to day of harvest. **CORN:** Use 10 to 20 lbs. per acre for corn earworm, Corn rootworm adults, European corn borer, fall armyworm, flea beetles, Japanese beetle, sap beetles and leafhoppers. For larvae in whorl (budworm damage) & foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. Use up to the day of harvest.

"Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations."

**CABBAGE, BROCCOLI, HEAD LETTUCE, CARROTS, POTATO, TOMATO, EGGPLANT:** Use 5 to 10 lbs. per acre for flea beetles and hornworm. Use 10 to 15 lbs. per acre for cabbage caterpillars, armyworms, & corn earworms. Avoid excessive applications. Some leaf injury may occur if treated plants are made when tender foliage is wet or in the presence of prolonged high humidity.

Do not apply within three days of harvest on Cabbage, Head Lettuce and other cole crops. For cucumber beetle, cutworm, diamond borer, squash bugs, and squash borer on cucumbers, melons, pumpkins and squash apply 10 lbs. per acre. For pepper, flea beetles 10 to 15 lbs. per acre.

**PEAS:** For the control of blister beetles and Japanese beetles use 20 lbs. per acre.

**NOTE:** Our recommendation for use of this product are based upon test procedures to be reliable. The use of this product by any person other than the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use. The results to be obtained must be based on directions or established safe practice. The buyer must assume a responsibility for the injury or damage caused by the use of this product in combination with other materials.