

Copper Sulfur Dust

THIS PACKAGE CONTAINS

ACTIVE INGREDIENTS

Copper (in copper oxide form) expressed as metallic 400

Sulfur 65.00

INERT INGREDIENTS 31.00

100.0

Boron from sodium borate pentahydrate 3.34

Read Entire Label Use in Strict Accordance With Label Directions and Cautions

ACCEPTED

CAUTION:

DIRECTIONS FOR USE

For the control of leaf spot. Make one application between June 20 and July 10.

At the rate of 15 lbs. per acre to supply $\frac{1}{2}$ lb. of boron per acre.

If additional material is needed apply Copper Sulfur Dust without boron as additional boron may injure the crop. Dust only when the foliage is dry.

NOTICE: Our recommendation for use of this product is based upon tests believed to be reliable. However, this product being beyond the control of the manufacturer, no guarantee expressed or implied is made as to the effects of such or the result to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage resulting from its misuse, as well as combination with other materials.

ANTIDOTE: Induce vomiting by tickling throat or an emetic or strong mustard water solution. Give 8 to 10 oz. In case of 1 to 2 tablespoonsfuls. Follow with large quantities of milk and honey. White oil eye drops and remove any foreign objects. Call a physician or take patient to hospital immediately.

POLYTRON *Surface Active Agent*

© 1958 E. I. du Pont de Nemours & Company

PRINTED IN U.S.A. 1958

EPA Reg. No. 1002