



NOVA TOXAPHENE 6E

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

See additional cautions on side panels.

ACTIVE INGREDIENTS:

Toxaphene*	60.25%
Petroleum Distillate	36.75%

INERT INGREDIENTS	3.00%
-------------------	-------

3% Emulsifiers (T-Mulz L and S, Thompson Hayward Chemical Co., (Kansas City, Kansas))

*Technical chlorinated camphene containing 67% to 69% chlorine.

F.P.A. Reg. No. 3624-95

EPA Est. 3624-KS-1

NET CONTENTS

GALLONS

NOVA PRODUCTS, INC.

Kansas City, Kansas 66119

DIRECTIONS

BEEF CATTLE, HORSES, SHEEP & GOATS: To control cattle lice, horn flies, cattle ticks (keds): Mix 5.33 pints of this concentrate with 100 gallons of water. (6 Oz. in 6 gallons will give the same concentration). Use a sprayer strong enough to part the hair of the animals and wet the skin. The animals may also be dipped. Treat when mites or lice appear and repeat treatment after 15 to 21 days if necessary. Calves should be sprayed sparingly but not dipped. (Do not apply to dairy cattle.) Do not spray within 30 days of slaughter. Do not use on dairy goats.

HOGS: To control lice and sarcoptic mange, mix 5.33 pints with 100 gallons of water. Treat when lice or mites appear and repeat treatment after 15 to 21 days if necessary. Do not dip pigs of less than weaning age. Do not treat animals within 45 days of slaughter.

GRASSHOPPERS: In corn or wheat fields: For young hoppers mix 1 quart with about 15 gallons of water or enough to cover 1 acre. Do not feed treated forage to dairy animals or animals being finished for slaughter. For mature grasshoppers increase dosage per acre from 1 quart in 15 gallons to 1 quart plus 10 ounces in 15 gallons of water. For small areas 1 tablespoonful to 1 gallon makes a suitable spray for about 200 sq. feet.

BAGWORMS ON EVERGREEN TREES: Mix 3 pints with 100 gallons of water (1.5 tablespoons with 1 gallon). Spray all the foliage thoroughly, adding a tablespoon of detergent to aid in the kill. Repeat as needed for control.

SOD WEBWORMS: For each 1,000 sq. feet of lawn mix 6 fluid ounces with 7 gallons of water and spray sod in June and again in August or September. Birds feeding in treated areas may be harmed.

ARMYWORMS—On Wheat, Oats, Barley, Sorghum and Rye, use 2 2/3 pints in 5 to 10 gallons of water per acre. See Caution Statement.

CUTWORMS—Variegated, Black and Subterranean Cutworms on Corn, Sorghum, use 2 2/3 pints in 10 gallons water per acre. Spray to give uniform application. For those working below ground, direct nozzles over row, apply in late evening. Cultivate dirt over row after treatment. See Caution Statement.

WARNING

May be fatal if swallowed. Do not breathe spray mist. Do not get on clothing or in the eyes. Do not contaminate food or feedstuffs. Do not apply to feed or forage crops to be fed to dairy animals or to animals being finished for slaughter. Do not dip animals under 1 month of age.

ENVIRONMENTAL CAUTION: This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditons favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

BEE CAUTION: This product is toxic to bees and should not be applied when bees are actively visiting the area.

CONTAINER DISPOSAL: Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

ACCEPTED WITH COMMENTS