NOPROKILL-FF

Contains 1% R BAYTEX

FOR USE BY QUALIFIED PEST CONTROL OPERATORS ONLY.

Ready To Use Residual Spray For Food Processing Plants, Restaurants, Hotels, Motels, Warehouses, Flour Mills, Breweries, Stores and Other Industrial Structures.

For residual control of cockroaches (waterbugs), ants, fleas (outside) mosquitoes, flies, crickets and wasps.

ACTIVE INGREDIENTS: 100% 0,0-Dimethyl 0- 4-(methylthio)-mtolyl phosphorothioate / 1.00% 2,2-Dichlorovinyl dimethyl phosphate* .46% Related Compound* .04% 98.50% Petroleum Distillate**

^R BAYTEX, registered trademark of ✓ U. S. Patent No. 3,042,703 Farbenfabriken Bayer A. G., Chemagro Canadian Patent No. 631,472 Corporation, licensee.

* Technical DDVP o ** Flash Point 176 F (TCC)

WARNING: Keep Out of the Reach of Children. See side panel for additional warnings.

NET CONTENTS GALLONS

Nova Products, Inc. Kansas City, Kansas 66119

DIRECTIONS

DOSAGE: For initial treatment or for heavy infestations apply at the rate of one gallon to 500 square feet. For subsequent treatments use one gallon to approximately 1,000 square feet of surface.

Meat Packing and Food Processing Plants; Apply as a coarse spray or with a paint brush to walls, ceilings, and other infestedareas. Use only in the inedible products areas of meat packing and poultry processing plants. Any exposed food products, equipment or packaging materials must be rémoved or covered before treatment. After treatment equipment and utensils must be thoroughly rinsed with water before resuming operations. Treat food processing plants early in the season before canning or other processing begins. Repeat applications, if necessary, when plant is not in operation.

ACCEPTED MAR 2 6 1968

UNDER THE FEDERAL INSECTICIDE TINGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER

TO ATTACHED COMMENTS.

USDA Reg. No. 3624-

Spraying unloading docks and outside areas where insects congregate will aid in preventing reinfestations. Good sanitation and proper disposal of wastes are essential in any effective insect control.

Inside Buildings:

Application should be made as a coarse spray wr with a paint brush to baseboards, cracks, crevices, and other insect hiding places. Also a apply to adjacent surfaces where insects will crawl when they come out of hiding. Heavy sprays may cause staining.

Exterior Use:

•,•

۰, ¢

.

٠

Apply as a coarse spray or with a paint brush to selected surfaces such as doors, screens, window frames, door sills, and areas of parches or patios where these insect alight or congregate. Apply to the nests of wasps after dark when all the insects have returned to the nest. Apply localized treatments to infestations of ants and fleas in outside areas where there is no vegetation. For ants, thoroughly wet hills and runways. Repeat as necessary to maintain control.

RESTRICTIONS

Do not use as a space spray. Avoid excessive wetting of asphalt tile, rubber, and plastic materials. Do not spray on plants. Do not spray in dairy barns or poultry houses. Do not use in areas wher e children or pets come in contact with treated surfaces. Remove fish or other pets from room before spraying.

WARNING.

May be fatal of swallowed, inhaled or absorbed through the skin. Rapidly absorbed through the skin. Do not get in eyes, on skin, or on clothing. Do not breathe spray mist. Wash thoroughly with soap and warm water after handling. Wash clothing in soap and hot water before reuse.

Avoid contamination of feed, food or food handling equipment. If food handling equipment becomes contaminated, clean thoroughly before reuse. KKX

ANTIDOTE: Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with stropine. NOPROKILL-FF

Inerts: None

.

•

. •

Nora Products, Inc., Kansas City, Kas.

•