

AQUEOUS FOOD PLANT FOGGING INSECTICIDE

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND
DOMESTIC ANIMALS
CAUTION: HARMFUL IF
SWALLOWED
DO NOT APPLY DIRECTLY TO
FOOD.
ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of takes streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its tabeling

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited PESTICIDE DISPOSAL—Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfiliapproved for pesticides.

CONTAINER DISPOSAL: Containers 30 gallons or more—a) reseal container and offer for reconditioning or b) triple rinse (or equivalent) and offer for recycling, reconditioning for disposal containers.

posal in approved landfill
Containers less than 30 gallons—As in bilabove
GENERAL!—Consult Federal State or local
disposal authorities for approved alternative
procedures

A Concentrated PYRENONE Micro Emulsion

Designed for Use in Edible Product Areas of Official Establishments Operating Under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs and in Other Food Processing Plants

Apply with Conventional Mechanical or Compressed Air Equipment Designed and Adjusted to Deliver a True Aerosol Mist

ACTIVE INGREDIENTS	
Pyrethrins) 5∾
*Piperonyl Butoxide, Technical	50% .
Petroleum Distillate	04
INERT INGREDIENTS92	500
100	0%

Equivalent to 4 (%) ib tylicarbity: (6 propylp peronyl) ether and to 1 (%) related compounds

PYRENONE Registered Trademark of Fairfield American Corporation

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

MANUFACTURED BY

NOVA PRODUCTS, INC.

Kansas City, Kansas 66119

The use of this product in to time periods when the plant in during treatment. All food; ment or thoroughly cleaned to

Close windows and doors a fogger or vaporizor adjusted to in diameter and 80% less thin cubic feet of space. Direct in area and behind obstructions the treated area and ventilate of

The above described applicate fruit files, mosquitoes, aliver scorpions, wasps, hornets, cit common stored product persuaw-toothed grain beetles, a Angoumois grain moths, Med moths, meal moths, and grain in

IN MUSHROOM PRODUCTLy, and Cecidomylida in growing of space. Fogger should be slightly upward over them cresults refrigeration should in rate of 1 ounce to 2000 cubic.

NE,

EPA Reg. No. 3624 172