

DIRECTIONS

1. For disinfecting and cleaning hard surfaces in meat and poultry production, such as in processing of meat, poultry, turkeys and equipment, dilute BIOGENE 1 ounce to 1 gallon of water (1:128). Wipe, mop or swab all surfaces with solution. Leave in contact 10 minutes. Rinse all surfaces coming in contact with food with potable water before re-use.
 2. For non-food disinfection of laying and broiler houses and hatcheries: Do not use in milking stalls, milking parlors or milk houses. Remove all animals, poultry and feeds from the premises. Remove all litter and manure from floors and other surfaces. Empty all feeding and watering troughs and appliances. Saturate all surfaces with a 1:128 solution. Ventilate buildings and other closed spaces. Do not house poultry, livestock or employ equipment until treatment has been absorbed or dried. All treated feed racks, waterers and fountains must be thoroughly scrubbed with detergents and rinsed with potable water prior to reuse.
 3. For general cleanup in food processing plants, patient rooms, surgery, nursing homes, all animal facilities, on walls, floors, hard surfaces, equipment. Use 1 ounce per 1 gallon of water (1:128). Cover surfaces thoroughly with solution and allow to remain in contact 10 minutes. For porous surfaces, increase concentration to 2 ounces per gallon. Rinse all surfaces coming in contact with food with potable water before re-use.
 4. For disinfecting and deodorizing garbage and trash receptacles, carts, hard surfaces and sick room equipment, use 1 ounce of BIOGENE to 1 gallon of water (1 part to 128 of water).
 5. BIOGENE is a prophylactic agent in preventing transmission of athlete's foot fungi. Prepare a solution of 1 ounce per 1 gallon of water; mop or swab this solution onto floors and other similar hard surfaces normally contacted by the feet in locker rooms, shower stalls, bathrooms and around swimming pools.
- BIOGENE is recommended for sanitizing in fogging operations to help reduce airborne bacteria. Fogging with 2 ounces of BIOGENE to 1 gallon of water 6 minutes per 1000 cubic of closed space will help reduce many airborne bacteria. Wait 2 hours after fogging is completed before re-entering the room.

F72

PHD6A

Net Contents _____ Gallons

OXFORD BIOGENE BACTERICIDAL DETERGENT

A concentrated one-step cleaner and powerful disinfectant-deodorant of proven effectiveness. The combination of multiple germicides and high active ingredients provides an effective one-step method of disinfecting and cleaning at a dilution of 1 ounce to 1 gallon of water (1:128).

FOR USE IN SANITATION IN MEAT AND POULTRY PRODUCTION - FOOD PROCESSING - HOSPITALS - ANIMAL AND POULTRY CARE - VETERINARY.

VIRICIDAL - SALMONELLACIDAL - STAPHYLOCIDAL - TUBERCULOCIDAL - FUNGICIDAL - PSEUDOMONACIDAL - CONDUCTIVE

Active Ingredients: Isopropanol, 12.00%; Potassium laurate, myristate, stearate compounds, 11.80%; Paratertiarybutylphenol, 3.45%; Ortho-phenylphenol, 3.45%; 4-Chloro-2-cyclopentylphenol, 1.87%; Tetrasodium ethylenediaminetetraacetate, 1.00%; Essential oils, 0.40%.

Inert Ingredients 66.03%

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food. Wash after using. Avoid inhalation of fogging mist.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

EPA Reg. No. 3635-128

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Do not reuse empty drum. Return to drum conditioner or destroy by perforating or crushing and burying in a safe place.

AOAC USE-DILUTION METHOD, 11th EDITION

Staphylococcus aureus	1:128
Salmonella choleraesuis	1:128
Pseudomonas aeruginosa (PRD-10)	1:128

AOAC OFFICIAL FUNGICIDAL TEST, 11th EDITION

Trichophyton interdigitale	1:128
----------------------------	-------

BIOGENE is a viricidal agent against lipophilic viruses, such as herpesviruses (strain HF) and influenza A2/Aichi/2/68 (Hong Kong) at 1:128 dilution.

BIOGENE is Salmonellacidal - effective against the organisms which cause pullorum disease, fowl typhoid, and paratyphoid.

AOAC USE-DILUTION METHOD, 11th EDITION

Salmonella pullorum	1:128
Salmonella typhimurium	1:128
Salmonella typhosa	1:128
Salmonella choleraesuis	1:128

BIOGENE is also effective against the following organisms using the AOAC USE-DILUTION METHOD, 11th EDITION.


Proteus vulgaris	1:128
Streptococcus mitis	1:128
Escherichia coli	1:128

AOAC TUBERCULOCIDAL TEST

Mycobacterium tuberculosis	1:128
----------------------------	-------

BIOGENE removes mildew stains from hard surfaces.

BIOGENE is conductive - may be used on conductive floorings as it does not interfere with electrical conductivity, as specified in the procedures in accordance with Paragraph 252, "Conductive Flooring of the National Board of Fire Underwriters", Specification No. 56.



Oxford Chemicals, Inc., P.O. Box 100, New York, N.Y. 10001

128