

For: Bakeries • Food Plants • Restaurants  
• Schools • Hotels • Motels • Office Build-  
ings • Hospitals and Food Handling Plants

ACTIVE INGREDIENTS:

\*+(5-Benzyl-3-furyl) methyl 2,  
2-dimethyl-3-(2-methylpropenyl)  
cyclopropanecarboxylate . . . . . 0.250%  
Related compounds . . . . . 0.034%  
d-trans Allethrin (allyl homolog  
of Cinerin I) . . . . . 0.250%  
Related compounds . . . . . 0.019%  
Aromatic petroleum  
hydrocarbons . . . . . 0.331%  
INERT INGREDIENTS. . . . . 99.116%

\*Cis/trans isomers ratio: Max. 30%

(±) cis. and min. 70% (±) trans.

+ U.S. Patent No. 3,465,007

**CAUTION KEEP OUT OF REACH  
OF CHILDREN**

Avoid contamination of feed and food-  
stuffs. Use in food processing plants only  
when plant is not in operation. Foods  
should be removed or covered during  
treatment. All food processing surfaces  
should be covered or thoroughly cleaned  
before using. In hospitals, patients should  
be removed from room prior to treat-  
ment. Room should be ventilated for two  
hours after spraying. Do not return pa-  
tients to room until after ventilation.

Remove pets, birds, and cover fish aqua-  
riums before spraying.

**WARNING:** Contents under pressure.  
Do not puncture. Do not use or store  
near heat or open flame. Do not expose  
container to temperatures above 120°F.  
Never throw container into fire or incin-  
erator. Harmful if swallowed.

Net Weight 16.5 Oz. (1 lb. 0.5 oz.)

Sold exclusively by EPA Reg. No. 3635-195  
EPA Est. 3635-GA-1

**OXFORD**  
CHEMICALS

A CONSOLIDATED FOODS COMPANY  
RESPONSIVE TO CONSUMER NEEDS

P.O. BOX 80202, ATLANTA, GA. 30341

shake well before each use. Remove pro-  
tective cap. Aim spray opening away from  
person. Push button to spray.

**For Houseflies, Mosquitoes, and Gnats:**  
Close all doors and windows. Point noz-  
zle upward. Direct the spray mist to all  
parts of the room, especially windows  
and other light sources which attract these  
insects. Fill the room with mist, then  
leave the treated area. Keep the room  
closed for at least 15 minutes. Ventilate  
the room when treatment is completed.

**Wasps:** Application should be made in  
late evening when insects are at rest.  
Spray liberally into hiding and breeding  
places, contacting as many insects as pos-  
sible. Repeat as necessary.

**For Crawling Insects:** Roaches, earwigs,  
grain mites, sowbugs, spiders and centi-  
pedes: Spray thoroughly into hiding plac-  
es such as sinks, under refrigerators, base-  
boards, and storage areas. Spray directly  
on insects where possible. Repeat as ne-  
cessary.

**Ants:** Spray trails, nests and points of  
entry. Spray on ants where possible. Re-  
peat as necessary.

**Fleas:** Protect dogs and cats from fleas  
by spraying their resting areas. Hold spray  
nozzle about 18 inches from area to be  
treated. Apply spray at the rate of two  
seconds per linear foot. Direct spray on  
resting areas and floor around these areas.  
Treat bedding the same way. Also spray  
entire inside surface of dog houses. Re-  
peat as necessary.

Also treat dogs and cats with a registered  
product before they re-enter the area just  
treated with this product.

A75

ISC-74

**ACCEPTED**  
APR 11 1975  
Under the Federal Insecticide,  
Fungicide, and Rodenticide  
Act, this product is registered for the use  
of . . . . . 3635-195