

OXFORD® SEPTICIDE

Disinfectant Bowl Cleaner

Staphylocidal – Fungicidal – Pseudomonacidal – Tuberculocidal

Septicide dissolves porous lime deposits and uric acid stains. Eliminates offensive odors.

When tested by the AOAC Use Dilution test, Septicide is effective against Staph. aureus, Salmonella choleraesuis and Pseudomonas aeruginosa (PRD-10). The product is effective against Mycobacterium tuberculosis when tested by the AOAC Tuberculocidal Activity Test, and is also effective against pathogenic fungi (Trichophyton interdigitale) by the AOAC Fungicidal Test. All tests done by the official methods of Analysis of the AOAC, Eleventh Edition, 1970.

DIRECTIONS

For Toilets— It is not necessary to remove the water from the bowl.

Use 1¼ ounces of Septicide to remove soil, rust, lime, scale and uric deposits, disinfect and deodorize.

Open flip cap, invert bottle and with light pressure, direct stream completely around underneath side of

Active Ingredients: Isopropanol, 10.00%; Hydrogen chloride, 8.19%; n-alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈) dimethyl benzyl ammonium chloride, 2.44%; n-alkyl (68% C₁₂, 32% C₁₄) dimethyl ethylbenzyl ammonium chloride, 2.44%; Phosphoric acid, 4.72%; bis (tributyltin) oxide, 1.00%; Essential oils, 0.50%

Inert Ingredients:70.71%

DANGER

KEEP OUT OF REACH OF CHILDREN

CORROSIVE: Causes eye and skin damage. Protect eyes and skin when handling this product. May be fatal if swallowed. Do not breathe vapor or fumes. Produces chemical burns. Do not get in eyes, on skin or on clothing. **FIRST AID: (Internal)**— Call a physician immediately. Drink a teaspoonful or more of magnesia or a cup of soapy water followed with milk of magnesia or raw egg whites. **(Eyes)**— Wash with water for 15 minutes. Get prompt medical attention. **(Skin)**— Wipe off the acid gently, immediately flood the surface with water, using soap freely, then cover with moist magnesia or baking soda.

Rinse empty container thoroughly with water and discard it.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

EPA Reg. No. 3635-199
EPA Est. No. 3635-GA-1



toilet bowl rim. Allow at least 10 minutes contact time for surface to be cleaned and disinfected. Wet bowl mop in solution and swab all surfaces including seat and lid. Rinse. Repeat application for cleaning heavily encrusted deposits.

On solid porcelain urinals— Place bowl mop over drain. Add 1¼ ounces of Septicide to one gallon of water. Follow the same procedure as for toilet bowls. Septicide will not damage solid porcelain china or vitreous ware.

Shower stalls— Glazed or other ceramic tile and grouting may be cleaned without damage. Use 1¼ ounces in a gallon of water, allow 10 minutes contact time for surface to be cleaned and disinfected, rinse thoroughly.

Septic tanks are not affected when used as directed.

Net Contents _____ Gallons

ACCEPTED


H75

OCT 2 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 3635-199

© 1972 CONSOLIDATED FOODS CORPORATION
ALL RIGHTS RESERVED

BC-26



P.O. BOX 80202, ATLANTA, GEORGIA 30341