

phila).
APPLICATION—cover all processing equipment, then by hand, electric foggers, air, or steam, spray until air is filled with fog
TIME—Fog or spray as long in advance of the next processing period as possible. Before next processing period flush all equipment with potable water.

CAUTION: INFLAMMABLE MIXTURE
 Do Not Spray in the Presence of
OPEN FLAMES OR PILE OF LIGHT
DESTROY CONTAINER WHEN EMPTY

MEAT PACKERS: Consult U.S.D.A. Meat Inspection Division Memorandum No. 52 and 214 for explicit restrictions.

GALLONS

USDA REG. NO. 3640-10



INSECTICIDE

A HIGH QUALITY CONTACT INSECTICIDE for the Control of Exposed Stages of Certain Insects Common to the Interiors of Food Processing Establishments

ACTIVE INGREDIENTS — 100.000%

Petroleum Distillate	99.187
N-octyl Bicycloheptene Dicarboximide*	42.
Piperonyl Butoxide**	.260
Pyrethrins	.130

*M.G.K. 264 Insecticide Synergist, U. S. Pat. Off. No. 2,476,512.
 **Consists of .208% (butyl carbityl) (6 propyl piperonyl) ether and .052% related compounds.

CAUTION:
 KEEP OUT OF THE REACH OF CHILDREN.
HARMFUL IF SWALLOWED
 Do not spray in the presence of exposed meats or other exposed foodstuffs. Remove edible products before spraying is started. Thoroughly clean premises—walls, benches, floors, and all fixtures with a suitable detergent solution after the completion of spraying operations, and before reuse of equipment.

Stearns Chemical Corp
 MANUFACTURING CHEMISTS
 MADISON, WISCONSIN

minutes. Sweep up dead or stupored insects and destroy them. Repeat treatment as often as necessary

TO KILL ROACHES: Spray liberally with force into all cracks and crevices in masonry and woodwork pipe openings in walls and floors, or wherever these insects are likely to be hiding, killing as many insects as possible. Repeat treatment when necessary.

TO KILL ACCESSIBLE STAGES OF CERTAIN STORED PRODUCT PESTS: For grain weevils, cadellies, grain moths, and grain mites spray at the rate of one gallon to 750 square feet of area to the point of runoff. Apply liberally to empty bins, chutes and elevator shafts. Apply to walls and floors, paying particular attention to cracks and crevices. Columns and ceilings in bins should also be sprayed, as should all spaces around and under equipment. Spray insects directly, but do not apply to foodstuffs. Repeat treatment as often as necessary.

TO CONTROL THE VINEGAR FLY (DROSOPHILA): Spray thoroughly incoming crated tomatoes or fruit, also unloading docks,—repeat as often as necessary.



[3640-10]

ACCEPTED
 JUN 11 1968
 U.S. DEPARTMENT OF AGRICULTURE
 FEDERAL BUREAU OF INVESTIGATION
 WASHINGTON, D.C.