

EXCELLENT - to keep insect control in...
phila.

APPLICATION - cover all processing equipment, then by hand,
electric foggers, air, or steam spray until air is filled with fog.

TIME - Fog or spray as long in advance of the
next processing period as possible. Before
next processing period flush all equipment
with potable water.

CAUTION: INFLAMMABLE MIXTURE
Do Not Spray in the Presence of
OPEN FLAME OR PILOT LIGHT
DESTROY CONTAINER WHEN EMPTY

CAUTION:
Foods should be removed or covered during
treatment. All food processing surfaces
should be covered during treatment or
thoroughly cleaned before using. When
using the product as a space spray in
these areas, apply only when plant is not
in operation.

MEAT PACKERS - Consult U.S.D.A. Meat Inspection Division
Memorandum No. 52 and 211 for explicit restrictions

GALLONS
U.S.D.A. Reg. No. 3640-11

INSECTICIDE

A HIGH QUALITY CONTACT INSECTICIDE

for the Control of Exposed Stages of Certain
Insects Common to the Interiors of Food Process-
ing Establishments

ACTIVE INGREDIENTS -- 100

Petroleum Distillate	98.100
N-octyl Bicycloheptane	
Dicarbonyl	1.000
Piperonyl Butoxide	.600
Pyrethrins	.300

*MGK 264 Insecticide Synergist,
U.S. Pat. Off. No. 2,476,512.
**Consists of .48% (butylcarbityl)
(6-propylpiperonyl) ether and .12%
related compounds

CAUTION
KEEP OUT OF THE REACH OF CHILDREN
HARMFUL IF SWALLOWED

Do not spray in the presence of exposed meats or other exposed
foodstuffs. Remove all products before spraying is started.
Thoroughly clean benches, walls, benches, floors, and all fixtures
with a suitable detergent solution after the completion of spray-
ing operations. And before reuse of equipment

Repeat treatment as often as necessary.

TO KILL ROACHES: Spray liberally with force in all cracks
and crevices in molding and woodwork, pipe openings in walls
and floors, or wherever these insects are likely to hide, hitting
as many insects as possible. Repeat treatment when necessary.

TO KILL ACCESSIBLE STAGES OF CERTAIN STORED
PRODUCT PESTS: For grain weevils, cadelles, grain moths, and
gran mites, spray at the rate of one gallon to 750 square feet of
area to the point of runoff. Apply liberally to empty bins,
chutes, and elevator shafts. Apply to walls and floors, paying
particular attention to cracks and crevices. Columns and ceilings
between posts should be sprayed, as should all spaces around
equipment. Spray insects directly, but do not apply
to products. Repeat treatment as often as necessary.

TO CONTROL THE VINEGAR FLY (DROSOPHILA) Spray
thoroughly surrounding related tomatoes or fruit, also including
dock, as often as necessary.

Stearns Chemical Corporation

MANUFACTURING CHEMISTS

MADISON 4 WISCONSIN

A.C. [unclear]
Mar - 11 - 1968
3640-11

Camerman's Note
Poor Copy