

# 2% SEVIN® BEAN DUST

**DIRECTIONS:** To control Mexican Bean Beetle apply at the rate of 25 to 30 pounds per acre. Make the first application when insects or egg masses on the underside of the leaves are first seen. Thorough application to the lower as well as the upper leaf surface is necessary for good results. Repeat applications at 7 to 10 day intervals as necessary for control. Do not treat vines that will be fed to dairy animals or animals being finished for slaughter.

**ACTIVE INGREDIENTS:**

Carbaryl (1-Naphthyl N-Methylcarbamate)	2.00%
Sulfur	25.00%

**INERT INGREDIENTS**

73.00%

100.00%

### NOTICE

The U. S. Food and Drug Administration has set tolerances for pesticide residues. Crops having residues in excess of their permitted tolerances should not be marketed and to do so may violate State law and will violate Federal law when sold in interstate commerce. Such crops should not be marketed without removal of such excess residues. Residue tests indicate that this dust can be applied at rates directed above to beans and potatoes on the day of harvest without exceeding residue tolerances.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

Do not reuse empty container. Destroy when empty.

### CAUTION: KEEP OUT OF REACH OF CHILDREN

**HARMFUL IF SWALLOWED OR INHALED.** Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs. Do not feed treated vines to meat or dairy animals.

EPA Reg. No. 3743-255

3743-255

