

PHONE 354-0111

NET OFFICE DRAWER 527

Southern AGRICULTURAL CHEMICAL CO. INC.
4-10-73
3743

2-6-20

NET WEIGHT
25 LBS.

CABBAGE DUST

For the Combined Control of Cabbage Worms & Aphids & Downey Mildew

ACTIVE INGREDIENTS:

Toxophene* (*Technical chlorinated camphene chlorine content 67-69%)	20.0%
INERT INGREDIENTS	72.0%
	100.0%

FOR USE ON

CABBAGE: For the control of imported cabbage worms, aphids, and downey mildew. Make 20 lb. per acre applications depending on size of plants. Repeat application when necessary. Do not use on cabbage after heads begin to form. Do not use before plants have become well established in field. Do not apply unless foliage is dry. This dust is recommended only when combined aphids and worm control and fungicide is necessary and is particularly effective where DDT-resistant cabbage worms are present.

WARNING: This dust is highly toxic and should be applied only with power equipment by experienced personnel.

WARNING: Toxic to fish and wildlife, do not contaminate streams and lakes, or ponds with this material. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Warning: Keep out of Reach of Children

NOTICE: Follow directions carefully. Timing and method of application, weather and crop condition, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given here with.

WARNING: Do not apply where there may be grazing by livestock. Application should not be made when wind drift would carry dust to food crops, pastures, domestic animals, or people adjacent to field being dusted. Keep all persons out of operating area or vicinity where they may be in danger of drift. Treated areas should not be re-entered until drifting insecticides and volatile residues have dissipated.

WARNING

May be fatal if swallowed or absorbed through skin — Rapidly absorbed through skin — inhalation or skin contact may, without symptoms, progressively increase susceptibility to Parathion poisoning.

If you suspect exposure see a doctor — do not breathe dust or vapor, wear a proper respirator — Do not get on skin, in eyes, on clothing — If spilled on skin immediately wash thoroughly with soap and water — Do not store near food or feed products.

Whenever handling or applying parathion wear clean rubber or rubber-dipped gloves and clean water-proof or freshly laundered protective clothing (coveralls, cap, etc), covering exposed surfaces. Clothing must not get wet through. Destroy and replace gloves fre-

quently. (Airplane pilots should wear a full-face canister-type mask.) For any handling indoors provide mechanical exhaust ventilation.

Bury spillage; clean up area with strong solution. Bathe promptly with soap and water after using or handling. Wash clothing with soap and hot water before re-use.

Do not reuse empty container. Destroy by burying or burning. Stay out of smoke.

Not for use or storage in or around the home.

ANTIDOTE

Atropine is antidotal — consult physician for emergency supply

FIRST AID TREATMENT

Call a physician at once in all cases of suspected parathion poisoning. Remove contaminated clothing and wash the skin clean with copious application of soap and water to remove all traces of parathion. Remove patient immediately from the area where parathion is present. If swallowed, induce vomiting by giving warm salty or soapy water. 2-Pam is also antidotal and may be administered in conjunction with atropine.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PATIENT

Physicians' Note: Warning symptoms include weakness, headache, tightness in the chest, blurred vision, non-reactive pin point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps. Treatment: Give atropine, preferably by injection, grains 1-100 two or three tablets at once and parenterally or orally every hour as required up to 30 tablets or until pupils dilate. Never give morphine. Clear chest by natural drainage. Artificial respiration and/or oxygen administration may be necessary. Observe patient continuously 48 hours. Report all patients to cholinesterase inhibitors with or without warning of increased susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.