

40 W TOXAPHENE

ACTIVE INGREDIENT:

Toxaphene* 40%

INERT INGREDIENTS 60%

100.0%

*Technical Chlorinated Camphene (Chlorine Content 67.69%)

TOMATOES: CUTWORMS — Apply to soil as needed after transplanting. Use 5 lbs. per acre in sufficient water to give coverage. Do not apply one day before harvest.

WARNING - Keep out of Reach of Children!

WARNING: May be fatal if swallowed. Do not breathe dust or spray mist. Do not get in eyes, on skin or clothing. Wash thoroughly after using. Do not store near feed or food products. This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not use on dairy animals. Do not feed treated crops to dairy animals or animals being finished for slaughter. Do not permit poultry to feed on treated crops. Dispose of used containers safely. Follow proper application schedules to eliminate the possibility of exceeding legally permissible residues.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

E. P. A. Reg. No. 3743-318

ACCEPTED	
JUN 19 1973	
U.S. DEPARTMENT OF AGRICULTURE	
PLANT INDUSTRY AND HOME ECONOMICS	
RESEARCH AND ADMINISTRATION SERVICE	
WASHINGTON, D.C. 20250	
SUBJECT TO ATTACHED COMMENTS	

3743-318