

ACCEPTED
 9/23/68
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 OF ECONOMIC POISON REGISTERED
 UNDER NO. 3997 SUBJECT



8 Fl. Oz.

SARCOPTIC MANGE TREATMENT

USDA Reg. No. 3997-1
 FOR TOPICAL USE ONLY

ACTIVE INGREDIENTS:

Gamma Isomer Benzene	
Hexachloride (from Lindane)	12.464%
Xylene	82.764%
INERT INGREDIENTS	4.772%
	<hr/>
	100.00%

USE: For the treatment of sarcoptic mange, lice, fleas and ticks on dogs, sheep, swine and beef cattle, as a spray or dip.

DANGER:
 KEEP OUT OF THE REACH OF CHILDREN.
 (See Inside Panel for Directions and Additional Cautions)

Arnold LABORATORIES, Cincinnati, Ohio 45209
 VETERINARY DIV. WESTERFIELD LABORATORIES, INC.

SARCOPTIC MANGE TREATMENT

As a spray or dip for the treatment of sarcoptic mange, lice, fleas and ticks on dogs, sheep, swine, and beef cattle follow the directions as given below. Repeat treatment after 2 or 3 weeks, if needed.

DOGS—Use 1 oz. in 4 gallons of water for sarcoptic mange mites, fleas and lice; 2 ozs. in 4 gallons of water for ticks.

BEEF CATTLE—Use 8 ozs. in 30 gallons of water. Do not exceed 0.03% on calves less than 6 months old.

SHEEP—As a spray, use 8 ozs. in 20 gallons of water; as a dip, use 8 ozs. in 40 gallons.

SWINE—Use 8 ozs. in 20 to 25 gallons of water as a spray or dip.

CAUTION: May be fatal if swallowed. Avoid contact with skin as may be absorbed. In case of contact, wash immediately with soap and water. Do not breathe spray mist.

WARNING: Do not DIP sheep, swine and beef cattle within 60 days nor SPRAY within 30 days of slaughter. Do not use on emaciated or lactating animals. Animals under 3 months should not be treated. Avoid contamination of animal feeds or water. Do not use on cats or humans. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter.

DANGER: KEEP OUT OF THE REACH OF CHILDREN.

Prescribed By

Arnold LABORATORIES, Cincinnati, Ohio 45209
 VETERINARY DIV.—WESTERFIELD LABORATORIES, INC.