

PHYSICAL OR CHEMICAL HAZARD:
Flammable. Contents under pressure.
Do not use near fire, sparks, or flame.
Do not puncture or incinerate container.
Exposure to temperatures
above 130° F. may cause bursting.

CAUTION: Avoid contact with skin.
Avoid breathing of vapor. Do not
remain in treated areas and ventilate
the area after treatment is completed.
Keep out of reach of children.

Do not spray directly against surfaces.
Do not use in any location where fish
or other aquatic or cold-blooded life
is kept. Avoid direct spray of food.

DISPOSAL: Do not reuse
empty container. Wrap
container and put in trash
collection.

Mfg. By:

ORB INDUSTRIES, INCORPORATED
52 Race Street, P.O. Box 1067
Upland, Pa. 19015
Tel: 215-TR 4-2537



TICK-TOX INSECTICIDE

No. 38

ACTIVE INGREDIENTS:

Pyrethrins	0.250%
Technical Piperonyl butoxide*	1.000%
Rotenone	0.128%
Other Cube Extractives	0.162%
Petroleum solvent	3.600%

*Equivalent to 0.8% (butylcarbitol) (6-propylpiperonyl) ether and 0.2% related compounds.

INERT INGREDIENTS TOTAL:

Isopropanol	15.00%
Methylene Chloride	15.00%
1, 1, 1 Trichloroethane	39.86%
Propane Isobutane	25.00%

EPA Est. No. 11-11-PA-1 EPA Reg. No. 10-39

Net Contents 12 oz. (340 g.)


CONTAINERS REFRIGERATE
DO NOT FREEZE
DO NOT HAND HEAT

DIRECTIONS

**FLIES, MOSQUITOES, GNATS,
SMALL FLYING MOTHS, WASPS:**
Close doors and windows to confine spray. Spray 4 to 5 seconds for a room 10 feet by 10 feet and larger rooms in proportion. Direct spray upward and in all directions to aid in dispersing spray. Keep doors and windows closed for 15 minutes after spraying. Repeat as required. Sweep up fallen insects.

**BIRDS, SILVERFISH & CERTAIN
OTHER CRAWLING INSECTS:**
Spray into cracks, crevices and other potential harborage to activate and drive out insect and then spray directly on insects present. Repeat treatments periodically as required.

FLEAS AND TICKS ON ANIMALS:
Spray animal directly so as to deposit a light mist. Comb or brush hair off not lay of hair to expose skin and allow spray to penetrate. Treat animal's back, underside and stomach, tail, back of neck and ears. Spray directly on ticks when possible. Do not get mist into animal's eyes or nose. Repeat weekly as required. Spray animal's sleeping quarters and bedding regularly as required to control any insects present and prevent reinfestation of animals.



BEST DOCUMENT AVAILABLE