

ORB #320 BAYGON[®] RESIDUAL INSECTICIDE

R Baygon is the registered tradename of Farbenfabriken
Bayer A.G. Chemagro Corporation Licensee

FOR INDUSTRIAL AND INSTITUTIONAL USE

KILLS: Roaches, Ants, Silverfish and
certain other crawling insects.

EPA Reg. No. 4077-67

ACTIVE INGREDIENTS

o-Isopropoxyphenyl methylcarbamate* 1.0%

Petroleum Distillate 84.0%

*U.S. Pat. #3111539

INERT INGREDIENTS

15.0%

**CAUTION: KEEP OUT OF REACH
OF CHILDREN. SEE ADDITIONAL
CAUTIONS ON RIGHT HAND PANEL.**

4077-67
ACCEPTED

NOV 13 1972

UNDER THE FEDERAL INSECTICIDE,
FUNGICIDE AND RODENTICIDE ACT
FOR E...
ED UNDER NO...
~~TO ATTACHED COMMENTS.~~

DIRECTIONS

This residual insecticide is intended for control of the insects listed in directions in such locations as storage areas, non-processing areas of food plants, restaurants, institutions, schools, homes, industrial plants, etc.

GENERAL: Apply with a low-pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off. Apply to accessible insect breeding sites and harbourages. Also treat adjacent exposed surfaces where insects may crawl when they come out of hiding. Repeat treatment as necessary.

COCKROACHES: Thoroughly treat the infested area. Spray baseboards, cracks and crevices; around window and door frames; the under surfaces of shelves, drawers, and work tables; behind and beneath cabinets, refrigerators, sinks and stoves; in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or nest.

ANTS: Apply to trails and places where ants enter premises such as window and door sills. Spray hills and runways outdoors.

EXPOSED STAGES OF THE FOLLOWING INSECTS: Cadellus, Cigarette Beetles, Drugstore Beetles, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Saw-toothed Grain Beetles and Weevils. Thoroughly treat floors and the walls of the infested area with attention to cracks, crevices, corners and perimeters. Sanitation and dust control are of primary importance. Fumigate or destroy infested products.

COCKROACHES: Thoroughly treat the infested area. Spray baseboards, cracks and crevices; around window and door frames; the under surfaces of shelves, drawers, and work tables; behind and beneath cabinets, refrigerators, sinks and stoves; in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or nest.

ANTS: Apply to trails and places where ants enter premises such as window and door sills. Spray hills and runways outdoors.

EXPOSED STAGES OF THE FOLLOWING INSECTS: Cadellae, Cigarette Beetles, Drugstore Beetles, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Saw-toothed Grain Beetles and Weevils. Thoroughly treat floors and the walls of the infested area with attention to cracks, crevices, corners and perimeters. Sanitation and dust control are of primary importance. Fumigate or destroy infested products.

SILVERFISH, FIREBRATS AND PAPERLICE: Apply to localized areas infested by these insects.

BROWN DOG TICKS: Spray sleeping quarter of pets, around baseboards, window and door frames, wall cracks and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not spray animals.

CLOVER MITES, CRICKETS, EARWIGS, MILLIPEDES, CENTIPEDES, SOWBUGS, SCORPIONS AND SPIDERS: Thoroughly spray the interior and exterior perimeter of the infested premises. For clover mites give special attention to window sills.

FLEAS, FLIES, GNATS, MOSQUITOES, PUNKIES and SANDFLIES (Outdoor Only): Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios and other areas where insects may enter.

FOR USE IN MEAT PACKING and FOOD PROCESSING PLANTS: Use only in the inedible product areas of meat packing plants, including those that are federally inspected. Any exposed food products, equipment or packaging materials must be removed or covered before spraying. Rinse all equipment with water before resuming operations. Treat food processing plants early in the season before canning or other processing begins. Repeat applications, if necessary, when plant is not in operation. Spraying unloading docks and outside areas where insects congregate will aid in preventing re-infestation. Good sanitation and proper disposal of wastes is essential in any effective insect control program.

CAUTIONS

Do not use as a space spray. Provide adequate ventilation of area being treated. Do not allow children to contact treated areas until surfaces are thoroughly dry. Do not store near food or food products. Do not use in commercial food preparation areas or in edible product areas of food processing plants. Avoid contamination of feed, food, food processing equipment and food handling surfaces. Do not apply to animals. Remove pets and cover fish bowls before spraying. Do not spray plants.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

May be harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin or on clothing. In case of contact, wash immediately with soap and warm water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Should illness occur, get prompt medical aid.

To Physician - Atropine Sulfate is antidotal.

Flammable: Do not use or store near fire or open flame.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place.