

Front Panel

Big Copy,
ONGARO'S WEED & FEED 20-5-5
Size: 20 lbs.
January 4, 1968



COVERS
5,000 SQ. FT. \$6.75
(100 X 50 FT.)

ONGARO'S
WEED & FEED

USDA REG. NO. 4185-354

INGREDIENT STATEMENT AS HERBICIDE:

ACTIVE INGREDIENTS:

Silvex (2-(2,4,5-Trichlorophenoxy) Propionic Acid)* 0.78% 1.4%

Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid** 0.62%

INERT INGREDIENTS Relative to Herbicide Content TOTAL 98.67%
100.0%

*Equivalent to 0.52% (2-(2,4,5-Trichlorophenoxy) Propionic acid).
**Equivalent to 0.52% 2,4-Dichlorophenoxyacetic acid (2,4-D).

CAUTION: KEEP OUT OF REACH OF CHILDREN AND PETS - SEE BACK PANEL FOR ADDITIONAL CAUTIONS

Gussets

ONGARO'S WEED & FEED

Back Panel

20 lbs. net weight


Covers 5,000 sq. ft.

ONGARO'S

WEED & FEED
20-5-5

GUARANTEED ANALYSIS:

TOTAL NITROGEN (N) 20.00%
6.00% Water Insoluble Nitrogen
AVAILABLE PHOSPHORIC ACID (P₂O₅) 5.00%
**SOLUBLE POTASH (K₂O) 5.00%
Total Available Primary Plant Food 30.00%
Potential Acidity Equivalent to 200 lbs. Calcium Carbonate per ton.
*In Michigan the heading reads, "MINIMUM GUARANTEED ANALYSIS."
**In states requiring, reads Water Soluble (K₂O).



Buy Copy
ONGARO'S WEED & FEED 20-5-5
Size: 20 lbs.

Back Panel (continued)

USDA REG. NO. 4185-354

INGREDIENT STATEMENT AS HERBICIDE:

ACTIVE INGREDIENTS:

Silvex (2-(2,4,5-Trichlorophenoxy) Propionic Acid) 0.75%

Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid ** 0.62%

INERT INGREDIENTS Relative to Herbicide Content

1.4%
TOTAL 98.6%
100.0%

*Equivalent to 0.52% (2-(2,4,5-Trichlorophenoxy) Propionic acid).
**Equivalent to 0.52% 2,4-Dichlorophenoxyacetic acid (2,4-D).

Manufactured For
ONGARO'S NURSERY
&
GARDEN CENTER, INC.
COLUMBUS, OHIO

Our premium Weed & Feed contains not one but two selected herbicides to rid your lawn of pesky weeds. These herbicides are scientifically impregnated in a high quality lawn food to give your lawn constant feeding over a longer period of time.

Spread it evenly over the lawn as directed - weeds will disappear and grass will take on a deep green color.

WEEDS CONTROLLED - Dandelion, ragweed, plantain, buckhorn, burdock, mustard, nettle, Jimson weed, lawn pennywort, pig weed, ground ivy and sow thistle.

CHICKWEED - is more easily controlled in very early spring. Several applications may be necessary. If applied during period of low temperature, there will be no immediate response from the lawn food. As the weather warms, the long lasting ingredients in the lawn food will be effective.

DIRECTIONS

Apply contents of this bag evenly over 5,000 sq. ft. of established lawn.
See spreader setting chart.

SOME THINGS TO REMEMBER -

Apply only when temperature is over 70 (except for chickweed).

Be sure there is plenty of moisture in the soil.

For quicker results, moisten grass lightly before applying.

Apply only to established lawns (lawns that have been mowed at least twice).

Apply evenly over entire area as directed.

If heavy rain occurs within 24 hours after application, second application may be necessary. (Wait at least one week before making second application).

Eq. No.
C-20-15-10-111-20-5-
Size: 20 lbs.



Back Page 1 (cont.) (end)

SOME HERBICIDES FOR USE -

- Do not mow for at least two days before applying.
- Do not mow for at least two days after applying.
- Do not water for at least 24 hours after applying.
- Do not sow seed for at least one week after applying.
- Do not apply on windy days.

CONDITIONS OF SALE. These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to plants and laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. The distributor, and the seller offer, and the buyer uses, this product subject to the understanding that the buyer assumes all such risks.

IMPORTANT: Use only on established lawns. Avoid drift of dust on vegetables, flowers, ornamental trees and shrubs, and other desirable plants. If some of the material comes in contact with desirable plants, it is recommended that plants be syringed thoroughly with water as soon as possible.

DO NOT store near other fertilizer, seed, insecticides or fungicides. Certain grasses such as Bent, St. Augustine, carpet grass and centipede are sensitive to herbicides and should not be treated.

CAUTION: Avoid prolonged or repeated inhalations of dust or contact with skin, eyes or clothing. Wash thoroughly after handling.

To protect fish and wildlife, don't use where runoff will contaminate streams, lakes or ponds. Drifting dusts, sprays, or vapors from this product may injure susceptible plants in the immediate vicinity.

WARRANTY AND LIMITATIONS OF DAMAGES. The distributor warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale; subject to the risks referred to therein. The distributor makes no other express or implied warranty, including any other express or implied warranty of **FITNESS** or of **MERCHANTABILITY**, and no agent of the distributor is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

SPREADER SETTING GUIDE

SCOTT	6	SEARS	6
LAWN BEAUTY	5	JACKSON	6
CENTRAL	5	AMC	5

These settings are approximate. Settings vary with model, age and condition of spreader.


January 4, 1968

1. Statement giving the sieve size of the granules and the method of incorporation of the herbicide:

The typical sieve analysis of the base or carrier is:

<u>Mesh</u>	<u>Cumulative Retained</u>
8 mesh tyler	Trace
14 mesh	10%
28 mesh	30%
through 28 mesh	70%

The herbicide is incorporated in the base by spraying a measured volume of 4 lb. concentrate into a weighed quantity of the base while this is being tumbled and mixed in a "Marian" ribbon mixer. The spray nozzles are designed to spray the charge evenly over the entire bed of material in about one minute. The charge is mixed an additional minute before discharge from the mixer to a screening and packagine unit.



Nutro

pre-emergent **CRABGRASS CONTROL**

14 - 3 -

GUARANTEED ANALYSIS*
 TOTAL NITROGEN (N) 14.00%
 8.7% Ammoniacal Nitrogen
 5.3% Urea Nitrogen
 3.0% Water Insoluble Nitrogen
 AVAILABLE PHOSPHORIC ACID (P₂O₅) 3.00%
 SOLUBLE POTASH (K₂O) 3.00%
 Total Available Primary Plant Food 20.00%
 Derived from—Primary Plant Food: Ammonium phosphate, urea, urea formylsulfide, sulfate of potash
 Potential acidity equivalent to 500 lbs. Calcium Carbonate per ton
 *In states requiring ready water soluble (RWS) ANALYSIS
 MSCA Reg. No. 4785 309
 Ingredient Statement as required
 ACTIVE INGREDIENT
 *Dimethyl Ester of Tetrachloroethylene Acid 3.85%
 INERT INGREDIENTS
 Relative to Herbicide Content 96.15%
 Total 100.00%
 *Use as a pre-emergent herbicide registered under U.S. Pat. No. 2,923,634
 Nutro Crabgrass Control Plus Weed Killer Turf Food is a pre-emergent herbicide which controls many species of crabgrass:
 1. Smooth Crabgrass (Dactyloctenium)
 2. Hair Crabgrass (Digitaria sanguinalis)
 3. Spiny Crabgrass (Eleusine indica)
 THIS PRODUCT ALSO CONTROLS THESE WEEDS:
 1. Annual Bluegrass 8. Florida Pussey
 2. Barnyard Grass 9. Goosegrass
 3. Bromus Paspalum 10. Green Foxtail
 4. Carpetweed 11. Hair Crabgrass
 5. Common Chickweed 12. Johnsongrass (from seed)
 6. Dandelion 13. Lambquarters
 14. Palmer Amaranth 21. Spring Pigweed
 15. Purslane 22. Spotted Spurge
 16. Prostrate Spurge 23. Storkgrass
 17. Redroot Pigweed 24. Texas Millet
 18. Redstem Filaree 25. Wildgrass
 19. Sesuvium 26. White Groundsweeper
 20. Smooth Crabgrass 27. Yellow Foxtail

Nutro Crabgrass Control Plus Weed Killer Turf Food not only prevents crabgrass and certain weeds, but it also stimulates the growth of grass and grass varieties not listed above. Nutro Crabgrass Control Plus Weed Killer Turf Food will prevent weeds and grass in flower beds.

DIRECTIONS
FOR USE ON LAWNS
 Distribute contents of this bag evenly over 2,500 sq. ft. of lawn area. Apply during early spring before crabgrass germinates. A good rule of thumb for the time to apply is to make the application prior to the time for applying cup its flowers.
 To assure even coverage, apply one half bag, then apply the other half, but disperse to the first application.
DO NOT MOW OR CULTIVATE AFTER APPLICATION HAS BEEN MADE

IMPORTANT: It is recommended that reseeding not be done the same year this product is applied. If it is possible to reseed 60 days after application, by increasing the amount of lawn seed sown by 50% to insure timely stands, this product can be applied to newly emerged grass after it reaches a height of one to two inches.
CAUTION: Do not contaminate feed or foodstuffs. Harmful if swallowed. Avoid contact with eyes. Wash after handling or using. Do not reuse con-

tainer, device when empty.
 Do not apply to newly seeded lawn until after green-up appears. Do not use on D Lawns. Do not store near seed, fertilizer or other pesticides.
 To control annual crabgrass and other late germinating grasses, seeds a late weed application at one half the recommended dosage.

DIRECTIONS
FOR ORNAMENTAL WEED CONTROL
 Nutro Crabgrass Control Plus Weed Killer Turf Food should be applied to newly sown weed seedling emergence or after the been cultivated and established weed. A good rule of thumb for timing the application is to make the application when Florida's temperature is 60°F and soil is still cold and weather is cool and humid.
 Plant beds should be evenly cultivated, transplanting or seeding. The surface of free of weed plants. Application should be to prevent weed seed germination.
 Contents of this bag are sufficient to treat 2,500 square feet of bed area. No supplementary treatments are necessary during growing season. **DO NOT** apply application. Control will include the annual weed and crabgrass control.

NUTRO CRABGRASS CONTROL PLUS WEED KILLER TURF FOOD IS REGISTERED AND SAFE FOR THESE ORNAMENTALS:
DECIDUOUS AND EVERGREEN TREES AND SHRUBS
 1. Aca 2. Abutilon 3. Ash 4. Asplenium

NET WT.

GOOD LAWNS MAKE GOOD
SMITH-DOUGLASS/DIVISION OF BORDE
COLUMBUS, OHIO 4

Purchase of this Nutro product entitles you to a free subscription to
NUTRO TURF & GARDEN GUIDE Magazine
 Printed in natural color with large pages for easy reading. Its seasonal hints and new ideas will save you time and money. If you would like a two-year FREE subscription, please complete this coupon.
 NAME _____ (please print)
 ADDRESS _____
 CITY _____ STATE _____ ZIP _____
 Mail to: Nutro Turf & Garden Guide, P.O. Box 600, Columbus, Ohio 43260

FREIGHT SHIPPING BAG
 Meeting requirements of Package 1348
 UNIFORM FREIGHT CLASSIFICATION
 Guaranteed by CHASE BAG CO

Nutro CRABGRASS CONTROL

Bug Copy
ONGARO'S WEED & FEED 20-5-5
Size: 20 lbs.
January 4, 1968

Front Panel

COVERS
5,000 SQ. FT. \$6.75
(100 X 50 FT.)

ONGARO'S
WEED & FEED

USDA REG. NO. 4185-354

INGREDIENT STATEMENT AS HERBICIDE:

ACTIVE INGREDIENTS:		1.4%
Silvex (2-(2,4,5-Trichlorophenoxy) Propionic Acid)*	0.78%	
Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid **	0.62%	
INERT INGREDIENTS Relative to Herbicide Content		98.6%
	TOTAL	100.0%

*Equivalent to 0.52% (2-(2,4,5-Trichlorophenoxy) Propionic acid).
**Equivalent to 0.52% 2,4-Dichlorophenoxyacetic acid (2,4-D).

CAUTION: KEEP OUT OF REACH OF CHILDREN AND PETS - SEE BACK PANEL FOR ADDITIONAL CAUTIONS

Gussets

ONGARO'S WEED & FEED

Back Panel

20 lbs. net weight


Covers 5,000 sq. ft.

ONGARO'S

WEED & FEED
20-5-5

GUARANTEED ANALYSIS:

TOTAL NITROGEN (N)	20.00%
6.00% Water Insoluble Nitrogen	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	5.00%
**SOLUBLE POTASH (K ₂ O)	5.00%
Total Available Primary Plant Food	30.00%
Potential Acidity Equivalent to 800 lbs. Calcium Carbonate per ton.	
*In Michigan the leading reads, "MINIMUM GUARANTEED ANALYSIS."	
**In states requiring, reads Water Soluble (K ₂ O).	



Bag Copy
ONGARO'S WEED & FEED 20-5-5
Size: 20 lbs.

Back Panel (continued)

USDA REG. NO. 4185-354

INGREDIENT STATEMENT AS HERBICIDE:

ACTIVE INGREDIENTS:	1.4%
Silvex (2-(2,4,5-Trichlorophenoxy) Propionic Acid)*	0.78%
Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid **	0.62%
INERT INGREDIENTS Relative to Herbicide Content	98.6%
TOTAL	100.0%

*Equivalent to 0.52% (2-(2,4,5-Trichlorophenoxy) Propionic acid).

**Equivalent to 0.52% 2,4-Dichlorophenoxyacetic acid (2,4-D).

Manufactured For
ONGARO'S NURSERY
&
GARDEN CENTER, INC.
COLUMBUS, OHIO

Our premium Weed & Feed contains not one but two selected herbicides to rid your lawn of pesky weeds. These herbicides are scientifically impregnated in a high quality lawn food to give your lawn constant feeding over a longer period of time.

Spread it evenly over the lawn as directed - weeds will disappear and grass will take on a deep green color.

WEEDS CONTROLLED - Dandelion, ragweed, plantain, buckhorn, burdock, mustard, nettle, jimson weed, lawn pennywort, pigweed, ground ivy and sow thistle.

CHICKWEED - is more easily controlled in very early spring. Several applications may be necessary. If applied during period of low temperatures, there will be no immediate response from the lawn food. As the weather warms, the long lasting ingredients in the lawn food will be effective.

DIRECTIONS

Apply contents of this bag evenly over 5,000 sq. ft. of established lawn.
See spreader setting chart.

SOME THINGS TO REMEMBER -

Apply only when temperature is over 70 (except for chickweed).

Be sure there is plenty of moisture in the soil.

For quicker results, moisten grass lightly before applying.

Apply only to established lawns (lawns that have been mowed at least twice).

Apply evenly over entire area as directed.

If heavy rain occurs within 24 hours after application, second application may be necessary. (Wait at least one week before making second application).

Bag Copy
GIGARO'S WEED & FEED 20-5-5
Size: 20 lbs.

Back Panel (continued)

SOME THINGS NOT TO DO --

- Do not mow for at least two days before applying.
- Do not mow for at least two days after applying.
- Do not water for at least 24 hours after applying.
- Do not sow seed for at least one week after applying.
- Do not apply on windy days.

CONDITIONS OF SALE. These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to plants and laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. The distributor, and the seller offer, and the buyer uses, this product subject to the understanding that the buyer assumes all such risks.

IMPORTANT: Use only on established lawns. Avoid drift of dust on vegetables, flowers, ornamental trees and shrubs, and other desirable plants. If some of the material comes in contact with desirable plants, it is recommended that plants be syringed thoroughly with water as soon as possible.

DO NOT store near other fertilizer, seed, insecticides or fungicides. Certain grasses such as Bent, St. Augustine, carpet grass and centipede are sensitive to herbicides and should not be treated.

CAUTION: Avoid prolonged or repeated inhalations of dust or contact with skin, eyes or clothing. Wash thoroughly after handling.

To protect fish and wildlife, don't use where runoff will contaminate streams, lakes or ponds. Drifting dusts, sprays, or vapors from this product may injure susceptible plants in the immediate vicinity.

WARRANTY AND LIMITATIONS OF DAMAGES. The distributor warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale; subject to the risks referred to therein. The distributor makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of the distributor is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

	SPREADER SETTINGS GUIDE		
SCOTT	6	SEARS	6
LAWN READER	5	JACOBS	6
CENTRAL	5	AM	5

These settings are approximate. Settings vary with model, age and condition of spreader.