

Front Panel

Bag Print (Paper)  
NUTRO WEED KILLER  
Size: 10 Lbs.  
May 9, 1969 - Rev. #4

TREATS  
5,000 SQ. FT.  
\$3.95

BORDEN  
TM  
Nutro  
(R)


DANDELION &

WEED KILLER

WILL NOT HARM GRASS

NET WT. 10 LBS.

INGREDIENTS



USDA Reg. No. 4185-404

ACTIVE INGREDIENTS:

Silvex (2-(2,4,5-Trichlorophenoxy) Propionic Acid)	3.01%
Propylene glycol butyl ether esters	1.21%
Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid	1.80%

INERT INGREDIENTS (Relative to herbicide content)

96.99%

Acid equivalents:

Silvex (2-(2,4,5-Trichlorophenoxy) Propionic Acid)	0.80%
2,4-Dichlorophenoxyacetic Acid	1.50%

CAUTION: KEEP OUT OF REACH OF CHILDREN AND PETS  
(See back panel for additional cautions)

TREATS  
5,000 SQ. FT.  
\$3.95

Nutro WEED KILLER

Gussets

BORDEN

Nutro WEED  
KILLER

Back Panel

NET WT.

BURDEN

10 LBS.

Nutro  
WEED KILLER

Back Panel (continued)

USDA Reg. No. 4185-404

Jul 9, 1969

USE	REGISTRATION
FILE	NO.
FOR	ACT
USE	REGISTERED
4185-404	

Bag Print  
NUTRO WEED KILLER  
Size: 10 Lbs.

ACTIVE INGREDIENTS:

Silvex (2-(2,4,5-Trichlorophenoxy) Propionic Acid)	3.01%
Propylene glycol butyl ether esters	1.21%
Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid	1.80%

INERT INGREDIENTS (Relative to herbicide content) 96.99%

Acid equivalents:

Silvex (2-(2,4,5-Trichlorophenoxy) Propionic Acid)	0.86%
2,4-Dichlorophenoxyacetic Acid	1.50%

BROADLEAF WEEDS CONTROLLED: Dandelion, ragweed, plantain, buckhorn, burdock, mustard, nettle, jimson weed, lawn pennywort, pig weed, ground ivy and sow thistle.

CHICKWEED AND OTHER SIMILAR WEEDS CONTROLLED: Common chickweed, mouse ear chickweed, little hop clover, bur clover, white clover, henbit, thyme-leaved speedwell, yarrow, knotweed and spotted spurge.

DIRECTIONS

Apply approximately 2 lbs. evenly over 1,000 sq. ft. (20' x 50') of lawn area. Contents of this bag will cover 5,000 sq. ft.

BROADLEAF WEEDS (Such as those listed above): For good results application should be made when the temperature is above 60° F., and there is adequate moisture in the ground so that broadleaf weeds are in a growing state. Chickweed may be controlled at the same time. Spring application is preferred, but late summer or early fall application is also recommended to destroy weeds that have developed from seed during spring and summer.

CHICKWEED: Application may be made whenever the temperature is above freezing. Chickweed can be controlled at lower temperatures than broadleaf weeds (a repeat application may be necessary).

IMPORTANT: Use only on established lawns. Avoid drift on vegetables, flowers, ornamentals, trees and shrubs and other desirable plants. If some of the material comes in contact with desirable plants, it is recommended that the plants be syringed thoroughly with water as soon as possible.

Do not store near fertilizer, seeds, insecticides or fungicides. Certain grasses such as Bent, St. Augustine, Carpet grass and Dichondra are sensitive to herbicides and should not be treated.

CAUTION: Avoid prolonged or repeated inhalations of dust or contact with skin, eyes or clothing. Wash thoroughly after handling. To protect fish and wildlife, don't use where runoff will contaminate streams, lakes or ponds. Vapors from this product may injure susceptible plants in the immediate vicinity.

CONDITIONS OF SALE. These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to plants and laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. BORDEN INC and the seller offer, and the buyer uses, this product subject to the understanding that the buyer assumes all such risks.

Front Panel

Bag Print (Paper)  
NUTRO WEED KILLER  
Size: 10 Lbs.  
May 9, 1969 - Rev. 4

TREATS  
5,000 SQ. FT.  
\$3.95

BORDEN  
TM  
Nutro  
(R)  
DANDELION &

WEED KILLER  
WILL NOT HARM GRASS

NET WT. 10 LBS.

INGREDIENTS

USDA Reg. No. 4185-404

ACTIVE INGREDIENTS:

Silvex (2-(2,4,5-Trichlorophenoxy) Propionic Acid)	3.01%
Propylene glycol butyl ether esters	1.21%
Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid	1.80%

INERT INGREDIENTS (Relative to herbicide content):

Acid equivalents:	96.99%
Silvex (2-(2,4,5-Trichlorophenoxy) Propionic Acid)	0.80%
2,4-Dichlorophenoxyacetic Acid	1.50%

CAUTION: KEEP OUT OF REACH OF CHILDREN AND PETS  
(See back panel for additional cautions)

TREATS  
5,000 SQ. FT.  
\$3.95

Nutro WEED KILLER

Gussets

BORDEN


Nutro WEED  
KILLER

Back Panel

NET WT.


BORDEN  
Nutro  
WEED KILLER

10 LBS.



Back Panel (continued)

USDA Reg. No 4185-404



Bag Print  
NUTRO WEED KILLER  
Size: 10 Lbs.



ACTIVE INGREDIENTS:

Silvex (2-(2,4,5-Trichlorophenoxy) Propionic Acid)	3.01%
Propylene glycol butyl ether esters	1.21%
Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid	1.80%

INERT INGREDIENTS (Relative to herbicide content) 96.99%

Acid equivalents:

Silvex (2-(2,4,5-Trichlorophenoxy) Propionic Acid)	0.86%
2,4-Dichlorophenoxyacetic Acid	1.50%

BROADLEAF WEEDS CONTROLLED: Dandelion, ragweed, plantain, buckhorn, burdock, mustard, nettle, jimson weed, lawn pennywort, pig weed, ground ivy and sow thistle.

CHICKWEED AND OTHER SIMILAR WEEDS CONTROLLED: Common chickweed, mouse ear chickweed, little hop clover, bur glover, white clover, henbit, thyme-leaved speedwell, yarrow, knotweed and spotted spurge.

DIRECTIONS

Apply approximately 2 lbs. evenly over 1,000 sq. ft. (20' x 50') of lawn area. Contents of this bag will cover 5,000 sq. ft.

BROADLEAF WEEDS (Such as those listed above): For good results application should be made when the temperature is above 60° F., and there is adequate moisture in the ground so that broadleaf weeds are in a growing state. Chickweed may be controlled at the same time. Spring application is preferred, but late summer or early fall application is also recommended to destroy weeds that have developed from seed during spring and summer.

CHICKWEED: Application may be made whenever the temperature is above freezing. Chickweed can be controlled at lower temperatures than broadleaf weeds (a repeat application may be necessary).

IMPORTANT: Use only on established lawns. Avoid drift on vegetables, flowers, ornamentals, trees and shrubs and other desirable plants. If some of the material comes in contact with desirable plants, it is recommended that the plants be syringed thoroughly with water as soon as possible.

Do not store near fertilizer, seeds, insecticides or fungicides. Certain grasses such as Bent, St. Augustine, Carpet grass and Dichondra are sensitive to herbicides and should not be treated.

CAUTION: Avoid prolonged or repeated inhalations of dust or contact with skin, eyes or clothing. Wash thoroughly after handling. To protect fish and wildlife, don't use where runoff will contaminate streams, lakes or ponds. Vapors from this product may injure susceptible plants in the immediate vicinity.

CONDITIONS OF SALE. These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to plants and laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. GORDEN INC and the seller offer, and the buyer uses, this product subject to the understanding that the buyer assumes all such risks.