

ACCEPTED

Aug 1, 1969

UNDER THE FEDERAL INSECTICIDE AND FUNGICIDE ACT FOR REGISTRATION UNDER THE POISON REGISTER 4581-151

the comments

..... Lbs. Net Weight

5% MALATHION DUST

ACTIVE INGREDIENT

*Malathion 5.0%

INERT INGREDIENTS 95.0%

TOTAL 100.0%

*O-O-dimethyl dithiophosphate of diethyl mercaptosuccinate

CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid prolonged breathing of dust. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Highly toxic to fish. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

USDA Registration No. 4581-151



AGRICULTURAL CHEMICALS

TACOMA, WASHINGTON

PHILADELPHIA, PA.

OAK BROOK, ILL.

BRYAN, TEXAS

FRESNO, CALIF.

MONTGOMERY, ALA.

DIRECTIONS FOR USE

5% Malathion Dust may be applied with air and ground equipment when air is calm. Application should be made covering the underside of the leaves as well as the top. The usual rate of application for the following crops is between 20 to 30 pounds per acre.

BEANS—For the control of Mexican bean beetle, aphids, leaf hoppers—make 2 or more applications as needed with no application later than 1 day before harvest.

PEAS—For the control of pea aphid, pea weevil on peas—apply to foliage during early bloom, repeat as necessary with no application later than 3 days before harvest.

POTATOES—For the control of aphids, leafhoppers—apply on foliage of infested plants and repeat as necessary.

BROCCOLI, CABBAGE, KALE, TURNIP, AND MUSTARD GREENS—For the control of aphids, imported cabbage worms, cabbage looper—apply on foliage and repeat as necessary with no application later than 3 days before harvest for broccoli and turnips and no later than 7 days before harvest for cabbage, kale, and mustard greens.

CATTLE—For the control of horn flies—apply on back and neck of the animal 3 tablespoons of 5% Malathion Dust per animal. Repeat at 10-14 day intervals. Do not apply to dairy cattle later than 5 hours before milking or during milking.

Do not contaminate feed, food containers, or water troughs.

Do not use in or near buildings; never re-enter. Stay out of smoke and fumes.

NOTICE

Pennwalt Corporation warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. Pennwalt Corporation makes no other express or implied warranty, including any other express or implied warranty of FITNESS or MERCHANTABILITY, and no agent of Pennwalt Corporation is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential damages such as loss of profits or value.

MADE IN U.S.A.

PRINTED IN U.S.A.

MALATHION DUST

	5.0%
	95.0%
TOTAL	100.0%

phosphate of diethyl mercaptosuccinate

CAUTION KEEP OUT OF REACH OF CHILDREN

Avoid prolonged or repeated contact with dust. Avoid contamination of feed and foodstuffs. Do not contaminate any body of water nor apply when crop area treated.

Highly toxic to bees exposed to direct contact. Protective information available from your Cooperative Agricultural Extension Service.

Registration No. 4581-151

PENNWALT
AGRICULTURAL CHEMICALS

PHILADELPHIA, PA.

FRESNO, CALIF.

MONTGOMERY, ALA.

air is calm. Application should be made covering the underside of the leaves as well as the top. The usual rate of application for the following crops is between 20 to 30 pounds per acre.

BEANS—For the control of Mexican bean beetle, aphids, leaf hoppers—make 2 or more applications as needed with no application later than 1 day before harvest.

PEAS—For the control of pea aphid, pea weevil on peas—apply to foliage during early bloom, repeat as necessary with no application later than 3 days before harvest.

POTATOES—For the control of aphids, leafhoppers—apply on foliage of infested plants and repeat as necessary.

BROCCOLI, CABBAGE, KALE, TURNIP, AND MUSTARD GREENS—For the control of aphids, imported cabbage worms, cabbage looper—apply on foliage and repeat as necessary with no application later than 3 days before harvest for broccoli and turnips and no later than 7 days before harvest for cabbage, kale, and mustard greens.

CATTLE—For the control of horn ^{flies} apply on back and neck of the animal 3 to 5 teaspoons of 5% Malathion ^{dust} per animal. Repeat at 10-14 day intervals. Do not apply to dairy cattle later than 5 hours before milking or during milking.

Do not contaminate feed, food containers, or water troughs. Do not burn or use in any containers; never reuse. Stay out of smoke and fumes.

NOTICE

Pennwalt Corporation warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. Pennwalt Corporation makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Pennwalt Corporation is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential damages such as loss of profits or values.

MADE IN U.S.A.

5-9979-05

PRINTED IN U.S.A.