DIRECTIONS

Use the amount of Toxaphene specified in 10-20 gallons of water per acre for ground sprayers or 2-10 gallons of water per acre for aircraft application.

ARMYWORMS AND CUTWORMS: For armyworms and climbing and surface feeding types of cutworms on Wheat, Oats, Barley, Sorghum, Rye, Corn and Alfalfa, use 2-2/3 pints per acre. Spray to give uniform application. Do not apply more than once after heads start to form on Sorghum. Do not feed treated forage, hay or corn ensilage to dairy animals or animals being finished for slaughter. Do not apply to forage to be sold commercially or shipped interstate. Do not ensile treated forage from Sorghum. Do not apply to grain sorghum within 4 weeks of harvest. There are no limitations on use of harvested grain.

FOR GRASSHOPPERS on Wheat, Oats, Barley, Sorghum, Rye, Corn and Alfalfa, as well as fence rows, use 2 to 2-2/3 pints per acre. For immature grasshoppers use low rate. Observe all cautions listed above.

CLOVER LEAF WEEVIL: For alfalfa in Missouri and Adjacent States, use 2-2/3 pints per acre. Apply in early Spring or after cutting before new growth is 4 inches tall, when leaf ragging on Alfalfa appears and damage is occurring from Weevil grubs. Do not feed treated forage to dairy animals or animals being finished for slaughter. Do not apply to forage to be sold commercially or shipped interstate.

BLISTER BEETLES: On soybeans, use 2 quarts per acre. Direct spray over row to give uniform coverage. Do not apply within 21 days of harvest. Do not make more than two applications after pods form. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

TOXAPHENE CROP CONTROL

(Emulsifiable)
Contains 6 Pounds Technical Toxaphene Per Gallon
For The Control of Grasshoppers,
Army Worms, and Cutworms


WARNING:

KEEP OUT OF REACH OF CHILDREN See Side Panel for Additional Warnings

ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Ca	m-
phene containing 67% to 69%	
Chlorine)	60%
Aromatic Petroleum Hydrocarbons.	34%
INERT INGREDIENTS:	6%_
Total	100%

COLORADO INTERNATIONAL CORP., COMMERCE CITY, COLO. 80022



TOXAPHENE CROP CONTROL

(Emulsifiable)
tains 6 Pounds Technical Toxaphene Per Gallon
For The Control of Grasshoppers,

Army Worms, and Cutworms

WARNING:

EEP OUT OF REACH OF CHILDREN
See Side Panel for Additional Warnings

FIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Ca	am-
phene containing 67% to 69%	
Chlorine)	60%
Aromatic Petroleum Hydrocarbons.	34%
RT INGREDIENTS:	6%
Total	100%

BAGWORMS ON EVERGREEN TREES: Mix 3 pints in 100 gallons water (1 tablespoon to 1 gallon water). Spray thoroughly all foliage. One tablespoon detergent added to spray mix will aid kill. Repeat as needed for control.

WARNING: May be fatal if swallowed. Do not breathe spray mist. Do not get on skin, clothing or in eyes. Wash thoroughly after using. Do not contaminate feed or food-stuffs.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

This product is poisonous to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on the label. EPA REG. NO. 4715-253

Net Contents___Gal.

ADO INTERNATIONAL CORP., COMMERCE CITY, COLO. 80022